



## GR Focus Review

# Rifted margins: Ductile deformation, boudinage, continentward-dipping normal faults and the role of the weak lower crust



Camille Clerc<sup>a,\*</sup>, Jean-Claude Ringenbach<sup>b</sup>, Laurent Jolivet<sup>c</sup>, Jean-François Ballard<sup>b</sup>

<sup>a</sup> LIVE, Université de la Nouvelle Calédonie, BPR4, 98851 Nouméa Cedex, France

<sup>b</sup> Total SA, CSTJF, Avenue Larribau, 64000 Pau, France

<sup>c</sup> ISTO-CNRS, Université d'Orléans, 1A rue de la Ferrollerie, 45071 Orleans Cedex 2, France

## ARTICLE INFO

## Article history:

Received 4 July 2016

Received in revised form 21 April 2017

Accepted 28 April 2017

Available online 12 May 2017

## Keywords:

Passive margin

Boudinage

Rifting

Lower crust

South China Sea

Atlantic

## ABSTRACT

The stunningly increased resolution of the deep crustal levels in recent industrial seismic profiles acquired along most of the world's rifted margins leads to the unraveling of an unexpected variety of structures. It provides unprecedented access to the processes occurring in the middle and lower continental crust. We present a series of so far unreleased profiles that allows the identification of various rift-related geological processes such as crustal boudinage, ductile shear and low-angle detachment faulting, and a rifting history that differs from the classical models of oceanward-dipping normal faults. The lower crust in rifted margins appears much more intensely deformed than usually represented. At the foot of both magma-rich and magma-poor margins, we observe clear indications of ductile deformation of the deep continental crust along large-scale shallow dipping shear zones. These shear zones generally show a top-to-the-continent sense of shear consistent with the activity of Continentward Dipping Normal Faults (CDNF) observed in the upper crust. This pattern is responsible for a migration of the deformation and associated sedimentation and/or volcanic activity toward the ocean. We discuss the origin of these CDNF and investigate their implications and the effect of sediment thermal blanketing on crustal rheology. In some cases, low-angle shear zones define an anastomosed pattern that delineates boudin-like structures. The maximum deformation is localized in the inter-boudin areas. The upper crust is intensely boudinaged and the highly deformed lower crust fills the inter-boudins underneath. The boudinage pattern controls the position and dip of upper crustal normal faults. We present some of the most striking examples from the margins of Uruguay, West Africa, South China Sea and Barents Sea, and discuss their implications for the time-temperature history of the margins.

© 2017 International Association for Gondwana Research. Published by Elsevier B.V. All rights reserved.

## Contents

1.	Introduction . . . . .	21
2.	Observations from a selection of industrial seismic reflection profiles. . . . .	23
2.1.	Uruguay magma-rich rifted margin. . . . .	23
2.1.1.	Observations. . . . .	23
2.1.2.	Interpretation . . . . .	23
2.2.	Southern Angola (Namibe) magma-poor margin. . . . .	24
2.2.1.	Observations. . . . .	24
2.2.2.	Interpretation . . . . .	25
2.3.	Gabon magma-poor rifted margin . . . . .	25
2.3.1.	Observations. . . . .	25
2.3.2.	Interpretation . . . . .	26
2.4.	South China Sea magma-poor margins . . . . .	26
2.4.1.	Observations. . . . .	26
2.4.2.	Interpretation . . . . .	27

\* Corresponding author.


E-mail address: [camille.clerc@univ-nc.nc](mailto:camille.clerc@univ-nc.nc) (C. Clerc).

2.5.	Barents sea magma-poor rift . . . . .	27
2.5.1.	Observations . . . . .	27
2.5.2.	Interpretations . . . . .	29
3.	Discussion . . . . .	30
3.1.	Ductile behavior of the continental crust . . . . .	30
3.2.	Crustal boudinage . . . . .	32
3.3.	Continentward Dipping Normal Faults (CDNF) . . . . .	34
3.3.1.	Origin of the CDNF. . . . .	34
4.	Conclusion . . . . .	36
	Acknowledgements . . . . .	36
	References . . . . .	37


**1. Introduction**

Rifted margins initially form as intra-continental rifts in response to continental divergence and evolve into passive margins when a new ocean forms. Studying the margins of oceans hence provides information on the mechanisms in action during continental rifting. Rifted margins are characterized by either an absence or an abundance of magmatic material. During rifting, the nature of the continental crust, its structural inheritance and its thermal state may lead to variable modes of crustal thinning. For scientific and economic motives, most of the world’s rifted margins have now been imaged by seismic reflection and improved technologies allow for ever-deeper penetration and better resolution. This influx of data makes it difficult to find an example of a “classical” model of rifted margin evolution (Levell et al., 2010). Most classical models of rifting were proposed on the basis of seismic profiles that do not show the resolution and penetration of more recent acquisitions. Numerical modeling suggests that the lower crust may in specific conditions behave as a ductile medium during its exhumation (Huismans and Beaumont, 2011, 2014) but these new images show structures in the lower crust, such as boudinage and shallow-dipping ductile shear zones that have been observed and considered in the last ten years in the Industrial World but have not been published yet. We show in this paper that the lower crust is often much weaker than classically postulated and this should lead us to reconsider rifting mechanisms in many cases.

Magma-poor rifted margins form as an answer to rifting and continental break-up without or with minimal syn-rift volcanism. Fig. 1 presents some of the many models proposed for the formation of magma-poor rifted margins. Some of them imply a ductile deformation of the lower crust (Huismans and Beaumont, 2011, 2014), with fluxes toward the ocean (Fig. 1a; Driscoll and Karner, 1998; Jolivet et al., 2015) or toward the continent (Fig. 1b, Brun and Beslier, 1996). Based on deep seismic data offshore United Kingdom, Reston (1988) proposes lower crust boudinage as an answer to extension (Fig. 1c). Based on their observations on the Tethyan margins preserved in the Alps and on the present-day Iberia Margin, Manatschal (2004), Péron-Pinvidic et al. (2007), Péron-Pinvidic and Manatschal (2008) and Mohn et al. (2012) proposed a model of three-layer continental crust where the brittle upper and lower crust are strongly decoupled by a ductile middle crust (Fig. 1d). The geometry of normal faults in this model, especially in the first stages, is influenced by the assumption that the lower crust is strong (intermediate or mafic) as in numerical models such as Lavier and Manatschal (2006), which leads to the formation of downward-concave normal faults for instance. Based on field observation in the pre-Pyrenean rifted margins, Clerc and Lagabrielle (2014), propose an alternative model implying boudinage of the upper crust allowed by a high geothermal regime and blanketing effect of the decoupled thick pre- and syn-rift sediments (Fig. 1e). Although the high temperature-low pressure metamorphism (up to 600 °C) recorded in these pre- and syn-rift sediment has not yet been reported from present-day passive margins this does not preclude their existence since drill holes are



**Fig. 1.** Some of the many models proposed for the behavior of the continental crust during the formation of rifted margins. (Source: modified after Reston, 2007).



**Fig. 2.** Variability in length of the margin, abundance of syn-rift deposits (sediments and/or volcanics), dip direction of the normal faults and general crustal behavior of the continental crust at rifted margins. Compilation from 15 rifted margins from this study and reproduced from Péron-Pinvidic and Manatschal (2008), Autin et al. (2010), Jolivet et al. (2015), Gernigon et al. (2014), Tsikalas et al. (2008), Untermeier et al. (2010), Osmundsen and Ebbing (2008), Clerc et al. (2015a), McDermott et al. (2015), Geoffroy et al. (2015), Gladczenko et al. (1998).


generally not deep enough and that such rocks are far out of target for industrial purposes.

The presence of a weak lower crust may favor an early necking and lead to an extensive spreading of the thinned continental crust (Huismans and Beaumont, 2011, 2014). For Brune et al. (2014) spreading of the continental crust is attributed to a rift migration maintained by sequential faulting in the brittle crust and lower crustal flow. For both Huismans and Beaumont (2011, 2014) and Brune et al. (2014), lower crustal viscosity is a key-parameter, which strongly depends on the composition of the lower crust, its initial thermal state and the extension rate during rifting. By contrast, the Newfoundland/Iberia conjugate margins show a wide zone of exhumed mantle resulting from low-angle normal faults rooted in a strong lower crust and upper mantle (Péron-Pinvidic and Manatschal, 2008).

The amount of extension measured from seismic profiles is generally insufficient to explain the amount of crustal thinning and subsidence (Ziegler, 1983; Burrus, 1984). This phenomenon, referred to as the extension discrepancy, is attributed to subseismic faulting (Marrett and Allmendinger, 1992; Crosby et al., 2008), polyphase faulting (Reston, 2005; Reston and McDermott, 2014), sequential faulting (Ranero and Pérez-Gussinyé, 2010) and/or to higher extension rates in the lower crust than in the upper crust, implying lateral flows of crustal material along shear zones (Sibuet, 1992; Davis and Kusznir, 2004; Egan and Meredith, 2007; Jolivet et al., 2015). Magma-rich margins are characterized by the presence of voluminous magmatic intrusion and extrusion contemporaneous with extensional tectonics (White and McKenzie, 1989; Skogseid et al., 1992). The main features of magma-rich rifted margins are: i) the presence of high-velocity seismic zones at the base of the crust; ii) a continental crust strongly intruded by sills and dikes and iii) covered with thick flood basalts and tuffs formation often imaged in seismic profiles as strongly reflective Seaward Dipping

Reflectors (SDR; Barton and White, 1997; Mutter et al., 1982; White et al., 1987; White and McKenzie, 1989).

The variety of tectonic models presented above is surpassed by the striking diversity in the geometry of the rifted margins observed worldwide. Several parameters such as the length of the margin, the thickness of the syn-rift sedimentary deposits, the behavior of the continental crust, the abundance of intrusive and extrusive magma and the vergence of normal faults differ significantly amongst margins (Fig. 2). Several Atlantic margins such as the Brazil/Angola conjugate margins present a strong necking zone followed seaward by a wide zone of strongly attenuated continental crust (c.a. 180 km wide), underlying a thick sag basin with an apparent lack of deformation (Contrucci et al., 2004; Moulin et al., 2005; Untermeier et al., 2010; Péron-Pinvidic et al., 2015). Whereas other margins such as the Southern Angola or Aden margins appear extremely short (e.g.: Autin et al., 2010). Such large variations in the width of the deformed domain, when not resulting from oblique rifting, call for drastically different rheological behavior already suggested by numerical modeling (Huismans and Beaumont, 2011, 2014), but never identified in nature. One then expects that a weak crust will have a tendency to spread more easily and lead to a wider margin, irrespective of other parameters such as tectonic heritage, which are probably more effective during the early stages of rifting. On this basis, it seems hardly possible to use a single model to understand and predict the geometry of all the rifted margins around the world. Although we are not yet able to propose a set of kinematic models as has already been done on some highly studied margins (Boillot et al., 1987; Whitmarsh et al., 1996; Pérez-Gussinyé et al., 2003; Péron-Pinvidic and Manatschal, 2008; Péron-Pinvidic et al., 2007), we present in this paper a selection of remarkable seismic reflection profiles that account for some geological processes rarely considered by the community of scientists working on rifted margins. We



**Fig. 3.** Location map for the 6 study areas. Basemap from <http://www.geomapp.org> and Ryan et al. (2009), repartition of the types of margins modified after Haupt et al. (2016). NAVP: North Atlantic Volcanic Province.

present: i) several observations in favor of a ductile behavior of the continental crust during the rifting of some rifted margins; ii) we introduce some hints in favor of tectonic boudinage at variable scales; iii) we present arguments in favor of the nucleation of rift-related low-angle normal faults; iv) we highlight the importance of the Continentward Dipping Normal Faults (CDNF) in the structuration of some rifted margins and discuss their implications.

## 2. Observations from a selection of industrial seismic reflection profiles

In this section, we present a selection of high quality seismic reflection data that illustrate some peculiar tectonic processes active during the formation of rifted margins (Fig. 3). These examples come from one magma-rich rifted margin (Uruguay) and four magma-poor margins (Gabon, Southern Angola, South China Sea and Barents Sea). For each seismic profile, we highlight the first order observations and then present our interpretations.

### 2.1. Uruguay magma-rich rifted margin

#### 2.1.1. Observations


The magma-rich rifted margin of Uruguay results from the break-up of Gondwana (Rabinowitz and LaBrecque, 1979; Gladczenko et al., 1997), which is *pro parte* contemporaneous with the emplacement of the Parana-Etendeka large igneous province (Gibson et al., 2006). In contrast with the magma-poor margins lying farther north, the Uruguay margin is devoid of post-rift evaporites but is characterized by thick accumulations, up to 8 km, of Seaward Dipping Reflectors (SDR)

interpreted as syn-rift volcanic formations (Stica et al., 2014; Franke et al., 2007; Clerc et al., 2015a).

In the lower crust of the Uruguay margin, Clerc et al. (2015a) identified two main groups of seismic reflectors (Fig. 4). A first group (A-reflectors) consists in long reflectors, continuous over tens of kilometers, especially abundant near the mantle-crust boundary. A second group consists of short reflectors (B-Reflectors) that are slightly curved and dipping toward the east and that tend to merge with the first group of reflectors (Fig. 4d). The concave-down geometry of the volcanics forming the SDR has sometimes been explained by subsidence models (Mutter et al., 1982; Hinz, 1981; Mutter, 1985), but alternatively, on-shore and offshore investigations have pointed out that SDR are syn-magmatic roll-over flexures controlled by major Continentward Dipping Normal Faults (Brooks and Nielsen, 1982; Planke et al., 2000; Gernigon et al., 2004, 2006; Geoffroy et al., 2001; Geoffroy, 2001, 2005; Mjelde et al., 2007; Stica et al., 2014; Pindell et al., 2014). Often masked by the thick overlying SDRs, those continentward-dipping normal faults are difficult to identify in seismic reflection. Fig. 5 presents an example of such a fault, clearly dipping toward the continent, and responsible for the oceanward tilt of a crustal block that supports a thick SDR wedge.

#### 2.1.2. Interpretation

Shear patterns observed at the base of the Uruguay margin are interpreted as the testimony of a ductile behavior of the base of the continental crust. The observed ductile shear could either be related to softening of the crust favored by high heat flow above the mantle plume, possibly underlined by younger intrusions, and/or to the deformation of syn-tectonic intrusions and underplated melt (White et al., 1987; White and McKenzie, 1989; Geoffroy, 2005).



**Fig. 4.** a) Regional map of the southern Atlantic Ocean after Gladczenko et al. (1997), Stica et al. (2014) and Soto et al. (2011), basemap from <http://www.geomapp.org> and Ryan et al. (2009). b) Magnetic anomaly map of the Uruguayan rifted margin after Maus et al. (2009). c) Interpretation of profile UR1 and d) close-up on the shear patterns at the base of the lower crust (Clerc et al., 2015a).

Source: Data courtesy of ION Geophysical.

Interestingly, the reflectors observed at the base of the continental crust draw a pattern that strongly resembles the one observed in ductile shear zones of various scales (Clerc et al., 2015a). The most characteristic feature calling for an analogy with ductile shear zones is the sigmoidal geometry of the small B-reflectors when approaching A-reflectors. The A-reflectors can be compared to the ductile shear planes whereas the B-reflectors are evocative of planes and objects deflected by the ductile deformation and parallelized to the shear bands. Together they draw the pattern of thick top-to-the-continent shear zones, consistent with the continentward dip of the normal faults that control the deposition of the SDR (Fig. 5). This observation suggests that the whole continental crust may have undergone a large-scale top-to-the-continent shearing. This further implies that, within the Ocean-Continent transition, lower crust and mantle have been exhumed by shearing during rifting, although the exact limits are difficult to position on the profiles. This behavior is similar in a first approach to the model recently proposed for the Gulf of Lion margin (Jolivet et al., 2015).

On the conjugate margins of Namibia (Fig. 1) and South Africa, several studies (Gladczenko et al., 1997; Mello et al., 2013; McDermott et al., 2015) describe comparable CDFN playing a first order role in the structuration of the margin and development of the SDR formations. The existence of SDR and of such antithetic CDFN on both sides of volcanic rifted domains (Geoffroy et al., 2015), though still debated (Becker et al., 2016), implies a symmetric disposition of structures that needs


to be discussed in term of mechanical behavior (see discussion in Section 3.3.1).

## 2.2. Southern Angola (Namibe) magma-poor margin

### 2.2.1. Observations

Limited southward by the Walvis Ridge, the Namibe margin is located at the southern end of the post-rift evaporites that cover most of the central segment of the South Atlantic (Jackson et al., 2000; Karner and Gambôa, 2007; Heine et al., 2013). The Namibe margin is extremely narrow in comparison with its conjugate Brazilian margin (Mohriak et al., 1995; Mohriak et al., 2008; Aslanian et al., 2009; Mohriak et al., 2012; Mello et al., 2013; Moulin et al., 2013). This asymmetry has been attributed to the development of upper versus lower plate boundary (Lister et al., 1986; Péron-Pinvidic et al., 2015), with the Namibe margin being the upper plate member (Brito et al., 2004). Because it is not covered by thick evaporite deposits, the Namibe margin allows for a fair imaging of the rift-related structures preserved in the upper continental crust.

Profile ANG1 (courtesy of ION Geophysical), presented in Fig. 6, is an 87 km-long profile that images the short Namibe margin, on the southwestern end of Angola. Numerous syn-rift basins are clearly imaged, characterized by oceanward-tilted fan reflectors.



**Fig. 5.** Profile UR2 and close-up on a SDR wedge and associated Continentward Dipping Normal Fault. Data courtesy of ION Geophysical.

**2.2.2. Interpretation**

As for volcanic rifted margins, most of the extension of the Namibe margin is accommodated by CDNF responsible for the deposition of seaward tilted syn-rift deposits (Fig. 6). But, unlike volcanic rifted margins, the Namibe margin is devoid of high impedance SDR and late-rift to post-rift magmatic activity is identified only at the western tip of the margin. According to the results of Moulin et al. (2013), there is no symmetry at the scale of the rifted domain since the Namibe margin is much shorter than its conjugate Brazilian margin.

This line demonstrates that magma-poor rifted margin can also be entirely controlled by CDNF. This implies tilting of the crustal blocks toward the ocean. This has strong implications in terms of sediments trapping because i) the deepest parts of the basins are located on the offshore side of each half-graben; and ii) the tilting of the sediments


adds up to the slope created by the subsidence of the margin whereas the dip of the normal faults is diminished by the subsidence of the margin (see discussion in Section 3.3).

**2.3. Gabon magma-poor rifted margin**


**2.3.1. Observations**

The Gabon margin was formed by the rifting apart of South America and Africa during the Late Jurassic and Early Cretaceous (Rabinowitz and LaBrecque, 1979). Onshore syn-rift sedimentary basins on the Gabon and its conjugate Brazilian margins indicate that rifting initiated in the Neocomian (i.e. 133 Ma; Teisserenc and Villemin, 1990; Magnavita et al., 1994). In contrast with the Angola margin, the Gabon margin presents a much thicker syn-rift sedimentary cover, and a thinner sag basin (Lentini et al., 2010; Unternehr et al., 2010; Péron-Pinvidic et al., 2015). We present three profiles shot offshore the Gabonese margin. 2D Profile GA1 and 3D profile GA2 (130 km-long) are parallel and very close to each other (Fig. 7). Profile GA1 is a 245 km-long and 24 km-deep (courtesy of Ion Geoventure). It shows the proximal margin from nearly undeformed continental crust in the east to well-developed oceanic crust in the west. Though the crustal reflectors are partly masked and blurred by thick post-rift evaporite, one can observe that the continental crust is thinned over > 150 km. On top of this thinned crust, the syn-rift sediments appear tilted toward the ocean. Profile GA2, shown as a line-drawing, (Fig. 7), 130 km-long and 16 km-deep, shows in more details the tilt of these sediments and reveals that the sedimentary reflectors are truncated/stopped oceanward along relatively low-angle continentward dipping reflectors. In Profile GA2, the continental crust is characterized by strong sub-horizontal reflectors presenting a low amplitude/long wavelength undulation.

GA3 (Fig. 8) is a 163 km-long 2D profile shot >100 km north of profiles GA1 and GA2, between the Koumaga and N’Komi Fracture zones (Mbina Mounguengui and Guiraud, 2009). It shares several characteristics with profiles 1 and 2: (i) two clear sets of oceanward-dipping thin sedimentary reflectors controlled by (ii) low-angle continentward-dipping normal faults that delineate (iii) an undulated top basement. A sharp bathymetric high is observed west of the shelfbreak. Although this profile is older and has a lower resolution close attention, It is worth noting that these features were hence already visible since the



**Fig. 6.** Map of West Africa showing the localisation of profiles GA1, GA2, GA3 and ANG1. Basemap from <http://www.geomapp.org> and Ryan et al. (2009). Profile ANG1 and interpretation.



**Fig. 7.** Profiles 1 and 2 from the Gabon rifted margin. Both profiles are parallel and roughly superposed. Approximate projection of profile GA2 on profile GA1 in indicated by the red box. See map on Fig. 6 for location.

Source: Data courtesy of ION Geophysical (profile GA1), Total, Marathon and Cobalt (profile GA2).

80ies-90ies but generally ignored because unpredicted by the models in force at that time.

### 2.3.2. Interpretation

Similar features can be observed at different scales and resolutions on profiles 1, 2 and 3. The syn-rift sediments are tilted toward the ocean ( $18^\circ$  to  $30^\circ$ ) thanks to the activity of CDNF. Identified by Meyers et al. (1996), these reflectors were interpreted as the gradational emergence of a proto-oceanic crust. But recent data show that these reflectors are distinct from conventional magma-related Seaward Dipping Reflectors (SRDs) by their low impedance and high frequency. Furthermore, there are no gravimetric or magnetic data indicating the existence of substantial syn-rift magmatic province in the area. The bathymetric high on profile GA3 is interpreted as a magmatic structure of limited extension, similar to other extrusive magmatic bodies already identified in the area (Jackson et al., 2000), but volcanism is not considered as a major process for the formation of the margin (Moulin et al., 2005). We therefore question, for the Gabon margin, the existence of a thick post-tectonic sag-basin resting on top of a long and flat detachment as generally represented along West African magma-poor rifted margins (Contrucci et al., 2004; Moulin et al., 2005; Unternehr et al., 2010; Péron-Pinvidic et al., 2015). Instead, we observe an in-sequence migration of extensional deformation (CDNF) and associated syn-tectonic sedimentation toward the ocean. Near the most distal part of profile GA3, late syn-rift sediments are associated to an increasing proportion of magmatic rocks (purple) that announces the forthcoming initiation of break-up and crustal spreading (Fig. 8). The most distal syn-rift sediments of profiles GA2 and GA3 rest


on a transitional domain that may either be constituted of exhumed mantle or proto-oceanic crust (Figs. 7 and 8). The nature of that strongly reflective contact is possibly tectonic. The thick syn-tectonic sediments are in turn locally covered by small and shallow sag-basins (yellow).

Below the seaward tilted syn-rift sediments, one can observe a continental crust characterized by flat reflectors that present a low amplitude (3 to 5 km) and long wavelength (10 to 50 km) undulation. On Profiles GA1 and GA2, strong intra-crustal horizontal reflectors at 11–13 km-depth, which is 2 to 5 km above the Moho, may correspond to the upper-lower crust limit visible in Martin et al. (2009) and identified by Dupré et al. (2011). These two crustal levels may be related to the two crustal levels “type I” and “type II” identified further south along the Angola Margin by Moulin et al. (2005) respectively interpreted as upper continental crust and lower (possibly intruded) continental crust. In agreement with Dupré et al. (2011), the lower crust imaged on profiles GA1, GA2 and GA3 appears irregularly shaped and presents lateral thickness variations along the direction of thinning and along the coast. We interpret these intracrustal undulations as upper crustal-scale boudins supporting oceanward-tilted syn-rift sediments above a sheared lower crust.

## 2.4. South China Sea magma-poor margins

### 2.4.1. Observations

The rifted margins of the South China Sea are devoid of salt and of SDR (Savva et al., 2014) and hence allow for a good imaging of the stretched continental crust.



**Fig. 8.** Seismic reflection profile GA3, line-drawing and interpretation across the Gabon margin, >100 km north of profiles GA1 & GA2. See map on Fig. 6 for location. Source: Data courtesy of Schlumberger Multisclient and Petrolin.

The triangular map-shape of the rifted domain with a SW pointing apex (Fig. 9), is indicative of a propagation opening (Courtilot, 1982; Huchon et al., 2001; Ding and Li, 2016). The “V” shape area is cored by a domain of Oligocene to Early Miocene oceanic crust. Because the South China Sea opened at the emplacement of the former long-lasting subduction zone of eastern Sundaland, it is characterized by a considerable amount of magmatic products uplifted and eroded since the Cretaceous (Taylor and Hayes, 1980, 1983; Schlüter et al., 1996; Pubellier et al., 2003; Wang et al., 2013). Rifting could have started from the orogenic collapse of the Cretaceous Yenshanian orogen (Chen et al., 2010) and mainly occurred from the Early Eocene to the Oligocene (Briais et al., 1993; Barckhausen and Roeser, 2004).

To the south west of the South China Sea, at the tip of the propagator, the rifted domain presents a ribbon morphology resembling the pattern of the Basin and Range province in Western US (Fig. 9). This pattern is generally considered as an extreme, pre-oceanisation, crustal thinning resulting in a boudinage of the whole continental crust (Nissen et al., 1995; Franke et al., 2014; Savva et al., 2013, 2014).

In Figs. 10 and 11, we present line-drawings of profiles SCS1 and SCS2. These 2 profiles are parallel, distant from one another by 60 km and are located across the Nam Con Son and East Natuna basins, at the very tip of the oceanic propagator (Fig. 9). The two profiles reveal a continental crust that is thinned and cut by normal faults over the whole length. On profile SCS1, little to undeformed 11 s-thick (tw) continental crust is attenuated to <5 s (tw). On profile SCS2, a 9 s-thick (tw) continental crust is thinned over >350 km, and continental break-up is probably achieved between two relatively symmetric margins.

Faults dips and dip directions are highly variable across and along the margins. (i) The northwestern half of profile SCS1 is characterized by dominant shallow-dipping CDFN whereas (ii) the dip direction of normal faults on its southern half and on profile SCS2 is much more variable with normal faults dipping both toward the ocean and toward the continent.

#### 2.4.2. Interpretation

Both profiles reveal a widely distributed thinning of the continental crust, which is strongly thinned and spread over >400 km due to two


episodes of rifting that occurred after the Yenshan orogen: the proto-South China Sea (now subducted beneath Borneo) and the South China Sea.

Profiles SCS1 and SCS2 highlight the importance of the along- and across-margin variability of the dip of normal faults. For Franke et al. (2014), this variability results from an alternating upper and lower plate configuration, but one could also invoke (i) the succession in time of several extensional events (Franke et al., 2014) or the unstable behavior of the continental crust in reason of its high thermal gradient (Cullen et al., 2010). On profile SCS2, the syn-tectonic sedimentary growth developed over the oceanward- and continentward-dipping normal faults appears diachronous. For instance, in Fig. 11b, the syn-rift sediments 1, in red, whose deposition is compatible with the activity of detachment 1, are overlapped by the syn-rift sediments 2, in orange, whose pattern is consistent with the activity of detachment 2. This change in the dip of the normal faults is responsible for the formation of crustal domes, commonly observed in the area. This example shows that most of the finite extension has been accommodated by low-angle normal faults in the upper crust and ductile thinning of the lower crust where normal faults root. The irregular topography of the Moho is a significant difference with the Basin and Range where the Moho is instead almost flat (Allmendinger et al., 1987). This difference is probably due to a larger finite extension and strain localization in a process that ultimately leads to breakup and formation of oceanic crust. Here, the upper crust is boudinaged during several episodes from Late Cretaceous to Miocene. Seismic refraction indicates that the large normal faults cutting through the brittle upper crust tend to flatten and root in a denser (6.5–6.7 km/s) and smoothly-contoured material interpreted as ductile lower crust (Pichot et al., 2014).

#### 2.5. Barents sea magma-poor rift

##### 2.5.1. Observations

Profiles BRT1, BRT2 and BRT3 lie in the southwestern Barents Sea (Figs. 12, 13), in an aborted branch of the post-Caledonian North Atlantic rift system (Riis et al., 1986; Breivik et al., 1998). The Barents




**Fig. 9.** Map of the South China Sea showing the main structural features, after Savva et al., 2013, 2014; Franke et al., 2014. Profiles SCS1 and SCS2 are located at the southwestern tip of the South China Sea propagator.

Sea opened over a part of the Caledonian orogeny, later inverted and eroded during the late Paleozoic (Gudlaugsson et al., 1998). The continental margin of the western Barents Sea developed in response to the separation of Eurasia and Greenland during the Paleogene (Talwani and Eldholm, 1977; Faleide et al., 1993). Following the orogenic collapse, the area, like the whole north Atlantic margin has undergone a long rifting history with peaks of activity during the Permian, the Jurassic, and the Early Cretaceous. Several basins, whose ages are progressively younger toward the West, are separated by basement highs (Loppa, Stappen, Veslemøy, Norsel, Senja). Gernigon et al. (2014) published a series of

crustal section based on deep seismic and potential field data modeling for the western Barents Sea (Fig. 12a). In this model, the Stappen and Loppa highs (Fig. 12a, b) represent massive and rigid crustal boudins, separated by a dominantly early Cretaceous basin, developed over thinned continental crust.

BRT1 is a 190 km-long profile across the Loppa and Veslemøy crustal high (Fig. 12c). Profiles BRT2 and BRT3, respectively 28,5 and 30,9 km-long, are from a 3D seismic block imaging the top of the Veslemøy high, down to 10 km depth. The area is characterized by a long low-angle reflector cutting through the sediments covering the top of the



**Fig. 10.** Profile SCS1, across the Nam Con Son basin, located ahead the propagator of the South China Sea. Source: Data courtesy of TGS.


Veslemøy high. The deep part of the low angle reflector is not clearly visible on seismic, nor is the basement (interpretation is possible down to the top Permian only). Seismic and gravimetric modeling show that the fault cuts the whole crust down to a dome-shaped Moho level at 25 km depth. Breivik et al. (1998) describe an inhomogeneous felsic to intermediate crystalline basement with seismic velocities of 6.3 km/s at the top to 6.6 km/s at the base. At the base of the crust, high velocities of 8.5 km/s are attributed to either mantle or eclogites.

3D seismic provides a good to fair imaging of the hanging wall down to 10 km, while the footwall is properly imaged down to 5–6 km. The shallow portion of the fault is a >1 km shear zone developed at the expense of the Triassic-Jurassic footwall and Cretaceous hanging wall. The hanging wall records several sets of conjugate normal faults; both antithetic and synthetic to the main shear zone, and cutting the sediments at low angle.


2.5.2. Interpretations

The necks between the crustal boudins have been characterized at low resolution in Gemigon et al. (2014). High-resolution 3D evidence the low-angle normal faults that were previously unresolved with the upper Mesozoic section. Between the Loppa and Stappen highs, large detachments appear to root in the very base of the crust and even cross through, down to the mantle, at 15 km to 25 km-depth. The

Veslemøy fault is very characteristic on this example; it affects the whole crust and is associated to a dome at Moho level (25 km instead of 32 km). The main activity of the fault is early Cretaceous; it has been reactivated to a lesser extent during the late Cretaceous Atlantic rifting and slightly rejuvenated after break-up. One important question is the original dip of this large-scale shear zone, was it initially steep and tilted in a later phase or was it formed with its present geometry? Thanks to 3D seismics, we were able to have a good 3D perception of the area and to pick and to measure the true dip of the associated minor conjugate of the normal faults. The contemporaneity of the conjugate faults is attested by their syn-sedimentary nature. We measured low dip values, ranging from 30° to 60° near the surface and down to ca. 14° at depth (Fig. 13a). By determining the bisector of the conjugate normal faults, it is possible to estimate the orientation of the major principal stress axis ( $\sigma_1$ ) at the time of faulting. Because  $\sigma_1$  is found vertical on Fig. 13b, and in opposition with what has been observed on other low angle normal faults (Reston and McDermott, 2011), we deduce that the faults have not been tilted after their formation. Furthermore, since these faults cut through the flat and unfolded Cretaceous sediments covering the top of the Veslemøy high, we can argue that these normal faults nucleated at low angle and are most probably not resulting from the inversion of pre-existing low angle thrust planes. This shows that the large-scale normal fault or extension shear zone



**Fig. 11.** A: Profile SCS2 across the Nam Con Son Basin, located 60 km northeast of profile SCS1. The seismic profile is available from Fig. 5A of Franke et al., 2014. B: Zoom on a crustal dome at the intersection of two antithetic detachments. Source: Data courtesy of TGS.



**Fig. 12.** a) Interpretation of the Barents margin redrawn after Gernigon et al. (2014). b) Localization map for profiles G, H, BRT1, BRT2 and BRT3, redrawn after Faleide et al. (1984) and Gernigon et al. (2014). c) Profile BRT1 across the Loppa and Veslemøy highs.

was formed with its present low dip like other detachments around the world: in the Basin and Range (Burchfiel et al., 1987), in the Cyclades (Mehl et al., 2005, 2007; Lacombe et al., 2013) or the Norwegian Caledonides (Osmundsen and Andersen, 2001). The possibility that normal faults can nucleate and be active at low-angle is a matter of debate. For some authors, nucleation is impossible and necessarily results from the reactivation of older fault planes and/or rotation of previous higher angle normal faults (Proffett, 1977). However, mechanical activation of low-angle normal faults is theoretically possible for material presenting very low friction coefficients (phyllosilicates, clays, talc, serpentinite...) and/or high fluid pressures (Famin et al., 2004; Collettini, 2011; Lecomte et al., 2011). This may be the case for the thick and reflective shear zone observed at depth on profiles BRT2 and BRT3, though it does not explain the low-angle normal faults observed on the surface. At a larger scale, interpretation of profile BRT1 reveals that the low angle normal faults affecting the top of the Veslemøy high relate to the fault system delimitating the northern rim of the Veslemøy crustal boudin. This indicates that the upper crust deformation is controlled by the boudinage with low-angle normal fault nucleating in the inter-boudin. This example thus shows a large-scale boudinage of the entire crust with a shallow-dipping extensional shear zone topping the boudins.


### 3. Discussion

#### 3.1. Ductile behavior of the continental crust

The rheology of the crust and its evolution during rifting is a matter of debate. According to numerous numerical approaches, ductile shear is suggested in the mid- to lower-crust during crustal thinning (Harry and Sawyer, 1992; Michon and Merle, 2003; Lavier and Manatschal, 2006; Van Avendonk et al., 2009; Huisman and Beaumont, 2011, 2014). However, in these models, most of the ductile deformation is localized in the middle crust whereas the lower crust is considered to be stronger and brittle (e.g. Van Avendonk et al., 2009), because resistant lower crust and upper mantle are chosen in the initial setups of these models. In contrast, other alternative quantitative models suggest a ductile behavior of the lower crust of some rifted margins following classical rheological stratification models where the lower crust is assumed to be made of intermediate to mafic material (Goetze and Evans, 1979; Le Pichon and Chamot-Rooke, 1991) or alternatively made of weaker quartz-felsic material (Hopper and Buck, 1996; Huisman and Beaumont, 2011, 2014 – Type 2; Jammes and Lavier, 2016). The examples shown in this paper suggest that for many margins the lower crust is weaker than assumed and accommodates a large part of extension by ductile shearing. Thanks to the exceptional quality of recent

seismic data, Clerc et al. (2015a) identified evidences for a rift-related ductile shear at the base of the Uruguay volcanic margin (Fig. 4). In the case of a volcanic margin such as the Uruguay one, the weakness of the crust could be related to the high heat flow related to the emerging plume and/or to the important syn-tectonic underplating and impregnation of the lower crust by magmatic melts (Franke et al., 2006; Hirsch et al., 2009). On the conjugate margin of Namibia/South Africa, high seismic velocities (up to >7 km/s) also point to a gabbroic composition of the lower crust (Fromm et al., 2015).

Other evidence of ductile deformation can now be interpreted at the base of the crust along the magma-poor rifted margins of Gabon and South China Sea. Along the Gabon margin, we observe 10 km to 50 km-long crustal-scale lenses that are reminiscent of boudins, so common on outcrops of ductilely deformed heterogeneous rocks. The base of the boudins, at a depth of 15 km to 12 km is separated from the Moho (17 km to 15 km-depth) by a 3 km to 5 km-thick lower crust characterized by long and sub-horizontal reflectors. This lower crustal layer is particularly evocative of a ductile behavior, especially in the inter-boudins necks where it bumps in between the stronger boudins (Fig. 14a and b). The geometry of the lower crustal layer in the inter-boudin and the attitude of syn-rift sediments strongly resemble, at a different scale, the shape of the salt diapirs observed at shallow levels on the same margins as well as the sediment packages deposited during gravity gliding on the rifted margin (Fig. 14c). This nearby analogy reinforces our perception that the lower part of the Gabon rifted margin behaves in a ductile way. The observed weakness of the base of the crust is possibly explained by a positive thermal anomaly during rifting suggested by Dupré et al. (2007, 2011). The origin of this thermal anomaly is still poorly constrained, but we suggest a potential effect of the sediment thickness. Indeed, crustal boudins are covered by thick syn-rift sedimentary deposits (4 to 8 km-thick) and the thick sedimentary cover is continuous, leaving no space for the heat to escape. The heating from below and insulation by sediment, or “blanketing effect”, are regarded as major issues for the thermal history of rifted margins (Levell et al., 2010; Clerc et al., 2015b, 2016; Lagabriele et al., 2016), and, as illustrated in Fig. 2, it is obvious that this effect is highly variable from a margin to another, depending on the sediment thickness and nature. Numerical modeling (e.g.: Karner, 1991; Lavier and Steckler, 1997; Theissen and Rüpke, 2010) confirms the blanketing effect of the sediments and resulting weakening of the lithosphere. On the other hand, the timing of rifting and associated sedimentation is another key factor, since the rapid input of sediment also affects upper mantle thermal structure and changes the buoyancy forces in ways that enhance the localization of strain and favor an early transition to narrow-rift mode (Lizarralde et al., 2007; Bialas and Buck, 2009). It is noteworthy that a ductile flow of the lower crust associated with a crustal-scale shear




**Fig. 13.** Profiles BRT2 (a) and BRT3 (b) across the Veslemøy crustal high (data courtesy of Schlumberger). Nucleation of low angle normal faults is attested from profile BRT2 (a), which is orthogonal to the faults strike (determined from 3D seismic block). b) The major principal stress  $\sigma_1$  is found vertical on BRT3 by determining the bisector of the conjugate antithetic normal faults, which attests for the absence of post-fault tilting. The low angle between the faults and the sedimentary bedding, and the tiny rotation of these series also confirms that the faults nucleate at low angle.

affecting the entire crust can lead to lateral juxtaposition of lower and upper crust. Once brought to shallow levels or even exhumed, a ductile lower crust is expected to behave in a brittle way and thus become undistinguishable in seismic reflection from an upper crust (e.g. Jolivet et al., 2015). This aspect should be kept in mind when attempting to

restore conjugate margins (cf. conservational vs. non-conservational models in Aslanian and Moulin, 2013).

In the South China Sea, tectonic boudinage and ductile deformation of the continental crust have already been proposed (Savva et al., 2013) and are in agreement with previous gravity modeling (Watts and



**Fig. 14.** a) interpretations of seismic profiles across the magma-poor margin of Gabon, a few tens of kilometers south of profile GA2. Courtesy of Total, Marathon and Cobalt. b) Comparison with similar structures observed at a completely different scale, in the rafted post-salt sediments. Localisation in (a). c) Comparison with a model of crustal boudinage, redrawn after Gartrell (1997).

Stewart, 1998) that suggests the existence of a mechanically weak zone underlying a wide part of the margin. Here, the ductile behavior is explained by the high heat flow resulting from the rapid inversion of a former magmatic arc (Nissen et al., 1995).

### 3.2. Crustal boudinage


Boudinage refers to an unstable behavior of a rheologically stratified medium subjected to extension (Goscombe et al., 2004). Stronger layers are deformed by periodical localized thinning leading to sausage-shaped structures separated by necks. These structures were named boudins by analogy with some sorts of French sausage (Lohest, 1909). Depending on kinematic boundary conditions, various shapes of boudins may form, that can be either symmetrical or asymmetrical (Goscombe et al., 2004). Boudins have been described on numerous outcrop scale examples but also at crustal scale (Fletcher and Hallet, 1983; Jolivet et al., 2004; Mehl et al., 2007) or even at lithospheric-

scale for the Basin and Range Province (Froidevaux, 1986; Ricard and Froidevaux, 1986). The wavelength of boudinage is highly dependent upon the rheological contrast between the boudinaged layer and the matrix and also its thickness like for all periodic instabilities. In the case of the Basin and Range, crustal-scale boudins are typically 30–50 km wide and lithospheric boudins show a 200 km periodicity (Ricard and Froidevaux, 1986; Ricard et al., 1987). Using the word “boudins” in the case of rifting and rifted margins is thus justified.

On several of the profiles presented above, boudinage appears as a recurrent deformation process accounting for the thinning of the continental crust at variable scales. On the margin of the Barents Sea, large (several tens to hundreds km), rigid crustal blocks such as the Loppa, Stappen, or Veslemøy blocks are interpreted as thick and poorly thinned crustal boudins (Gernigon et al., 2014). These crustal blocks are separated by inter-boudins characterized by thinned and sheared crust covered by thick syn-tectonics sediments. Similar features are described by Savva et al. (2013) in the South China Sea, where thin (ca. 2 to 4 s-

TWT) and elongated (100 km to 200 km-long) crustal boudins are separated by thick sedimentary basins floored by exhumed mantle peridotites. On a smaller scale, but still with comparable geometries, the distal part of the Gabon margin reveals the existence of 10 km to 50 km-long boudins of upper crust (Figs. 7, 8, 14). There, a network of numerous sub-horizontal reflectors in the lower crust delimitate crustal boudins, tapering at each end in an inter-boudin area controlled by low-angle ( $10^\circ$  to  $15^\circ$ ) CDNF. Similar anastomosed low-angle reflectors identified in the lower crust offshore Britain were interpreted as semi-brittle shear zones (Reston, 1988).

The boudin geometry distinguishes from the classical block tilting geometry described on rifted margins by (i) the shallow dip of normal faults and (ii) the lenticular or phacoidal geometry of the crustal blocks. Jolivet et al. (2004) and Mehl et al. (2007) highlighted the role of boudinage as an initial localizing factor by inducing the formation, near the brittle-ductile transition, of ductile shear zones at the edges of boudins. The inter-boudin necks tend to localize strain, which leads to local stress concentration and to initiation of shear zones. The shear zones are then progressively exhumed and replaced by shallow dipping cataclastic shear zones when they reach the brittle field (Jolivet et al.,



**Fig. 15.** a) Sand model uniformly stretched by downslope planar normal faults, redrawn after Vendeville, 1987. b) Sandbox experiment showing the progression of deformation sequence through an extensional wedge, photograph and redrawn from Xiao et al., 1991 and Yin, 1994. The locus of active synthetic normal faulting moves from the back to the front of the wedge (from right to left on the photograph). Comparison with profile ANG1 (c) suggests the existence a major shear at the base of the continental crust.

2004, 2010). This process leads to a final geometry consisting in boudins of variable size, separated by low angle shear zones rooted in inter-boudin necks, as exemplified on Fig. 12.


The observation of large-scale boudins in the extended crust of rifted margins calls for mechanical instabilities during rifting because of the stratification of the lithosphere, and more specifically the crust. The examples of the South China Sea or the Gabon magma-poor margins thus show a significant contrast of resistance between the upper and lower crust. The upper crust is periodically boudinaged above the weak lower crust and shallow-dipping extensional shear zones and normal faults form along the upper surface of these boudins. The exaggeration of this process leads to a boudinage of the whole crust (Barents Sea), possibly above a weak base of the crust or even above a hot and weak mantle.

### 3.3. Continentward Dipping Normal Faults (CDNF)


#### 3.3.1. Origin of the CNDF

The dataset we selected for this study, after browsing most of the world's margins, is remarkable for the important proportion of margins controlled by CNDF, which provides us with a rather unexpected perception of the magma-poor rifted margins, traditionally represented with

oceanward-dipping normal faults. The Namibe and Gabon margins are particularly striking with the whole margins being entirely controlled by CNDF. The exceptional quality of the data on profiles GA2, allows us to determine that the CNDF evolved in a sequence migrating toward the ocean, entailing with it an oceanward migration of depocenters (Fig. 7). Syntectonic sedimentation clearly results in a series of oceanward-growing and tilting sedimentary wedges, younging toward the ocean. The observation of CNDF along the Gabon margin is in accordance with previous observations (Ranero and Pérez-Gussinyé, 2010; Skogseid, 2014) and numerical models (Brune et al., 2014) but is also in relative contradiction with the description of the West African non-magmatic margins, often represented with thick and wide sag basin (Contrucci et al., 2004; Moulin et al., 2005) that is described as “subsiding vertically without differential tilting” (Aslanian et al., 2009) resting on a “highly thinned, little faulted continental crust” (Unternehrl et al., 2010). For us, interpretation of profiles GA1, GA2 and GA3 indicates that most of the sedimentary basins resting on top of the highly thinned continental crust are actually syn-rift and migrate outboard until break up. This implies that improved seismic images may lead us to reconsider the existence or extension of some of the sag basins interpreted on the thinned crust of several rifted margins.



**Fig. 16.** Possible scenario for the formation of Continentward Dipping Normal Faults (CNDF). a) Formation of CNDF by inversion of inherited structures (former thrusts), modified after Raillard et al. (1998) and Sapin et al. (2012); the formation of CNDF can also be controlled by the existence of top-to-the-continent shear at the base of the crust. This shear could result either from (b) a basal drag of the mantle due to mantle drift; or (c) in the case of volcanic rifted margins to the collapse of a topographic anomaly above an emerging plume.




**Fig. 17.** Implication of fault dip on the tilt of the crustal blocks. On CDNF-controlled margins, bloc-related tilting adds up to the subsidence-related tilting. Whereas on margins controlled by oceanward-dipping normal faults, block tilting (toward the continent) is opposed to subsidence-related tilting (toward the ocean).

Equivalents of the Continentward Dipping Normal Faults of present day margins can be observed in the Jurassic distal margins exhumed in the Alps. They are the Margna fault in the Malenco area (Müntener and Hermann, 2001) and the Pogallo shear zone in the Ivrea zone (Handy and Zingg, 1991). In these localities, Mohn et al. (2012) described a necking zone characterized by an interplay of converging normal faults rooting in the ductile middle crust. These faults, dipping continentward in the

lower crust (Pogallo fault) and oceanward in the upper crust (Grosina detachment), lead to the nearly complete elimination of the ductile middle crust. Once necked and sufficiently thinned, the crust becomes brittle which allows the faults to cut through, down to the mantle and exhumate it at the seafloor.

These processes, very well described in the Alps, seem to apply also fairly well to the Iberian margin (Mohn et al., 2012; Manatschal, 2004; Péron-Pinvidic and Manatschal, 2008). However, considering the important variability of the nature of the crust, the variability of its thermal state, and the diversity of extension rates, we may question the universality of this model. e.g.: Along the Cretaceous passive margins in the Pyrenean range, recent work revealed the existence upper crust boudinage (Clerc and Lagabrielle, 2014; Clerc et al., 2016; Corre et al., 2016; Teixell et al., 2016) associated to strong thermal anomaly recorded in the pre- and syn-rift deposits (Clerc et al., 2015b; Lagabrielle et al., 2016). Similarly, in the South China Sea, Savva et al. (2013) described a rift related crustal deformation widely distributed over numerous lower crustal detachments indicating a rather delocalized and ductile deformation of the crust. For Savva et al. (2013), these attributes suggest that the lithosphere was possibly warmer or weaker during the rifting of the South China Sea. Our observations in the same regions call for a similar interpretation.

Sandbox experiments by Faugeres and Brun (1984) and Faugeres (1985) revealed the relationship between the development of asymmetrical block tilting and the presence of a strong shear stress at the base of the brittle layer (Fig. 15). In these experiments, the dip direction of a normal fault is directly controlled by 2 parameters: (i) the inheritance of pre-existing tectonic discontinuities, and (ii) the sense of shear at the basal discontinuity in which the normal fault is rooted. Faugeres and Brun (1984) show that the normal faults are always synthetic to the basal shear (Fig. 15). It should be noticed here that the dip of normal faults rooting within shallow-dipping extensional shear zones along the upper surface of crustal-scale boudins mentioned in the previous section fully respect these characteristics. In an application of these models to rifted margins, Brun et al. (1985) describe an interplay between two systems



**Fig. 18.** Schematic representation of the volcanic rifted margins and ductile-type magma-poor rifted margin.

leading to the necking of the continental crust: (i) one corresponds to the upward migration of the Moho and affects the base of the crust with a top-to-the-continent sense of shear leading to the formation of CDNF; (ii) the other one corresponds to the subsidence of the margin, and leads to the tectonic collapse and gravity-driven décollement responsible for the formation of oceanward-dipping normal faults. By adding a layer of ductile silicone at the base of the sand layers, Vendeville (1987) observed more variable dip directions of the normal fault and the formation of horst and graben which could be attributed to the decoupling effect of the ductile layer. A looser transmission of the shear stress to the base of the brittle layer would hence lead to a more random organization of the dip of normal faults. It is tempting to compare this later result to the case of the South China Sea, which is known for its hot and thus weak and ductile lithosphere and where we observe an important along- and across-margin variability of the dip of normal faults (Figs. 10 and 11).

Therefore, in the case of the magma-poor rifted margins of the southern Atlantic (Gabon, Southern Angola) we may envision that the continentward dip of the normal faults could be either due to a structural inheritance (Fig. 16a) or to a large-scale shear at the base of the crust due to the upward migration of the Moho during necking. In the second situation simple shear then localizes along the Moho and controls the formation of CDNF in a symmetrical fashion like in the Southern Atlantic (Uruguay and Namibia conjugate margins; Figs. 2; 16b and c). Along the volcanic margin of Uruguay, this second hypothesis of a crustal shear at the origin of the formation of CDNF is indeed attested by the top-to-the-continent ductile fabric observed in the middle and lower crust (Clerc et al., 2015a). The shear pattern in the lower crust and the normal faults in the upper crust of the Uruguay margin show a component of simple shear distributed across the entire continental crust, thus suggesting a relative motion of the mantle with respect to the continental crust. When considering the conjugate southern Namibian and South African margins (Gładczenko et al., 1997; Blaich et al., 2013; Mello et al., 2013) we note a relative symmetry of the rift system since CDNF control the formation of SDR on both sides of the conjugate volcanic margins (Fig. 2).

A model supporting both (i) the top-to-the-continent basal shear and (ii) the existence of CDNF on each side of the rifted domain is also compatible with a gravitational collapse of a plume head-related topographic bulge for the volcanic margins (Fig. 16c). For instance, Griffiths and Campbell (1991) predict that the rise of a surface uplift around 400 km to 600 km wide can be expected above a 500 km radius plume head. We suggest that such a topography, in conjunction with the extension responsible for the opening of the Atlantic rift system could lead to a gravity collapse of the bulge resulting in a top-to-the-continent shear sense on each side of the rift. Lubrication of the base of the crust by intrusive and underplated magmas could weaken the crust and favor its collapse.

Because of the development of CDNF the crustal blocks are tilted toward the ocean. Later the rifted margin subsidence increases the dip. The resulting dip of the pre- and syn-rift sediments deposited on margins controlled by CDNF hence presents higher absolute tilt values than those deposited on margins controlled by oceanward-dipping normal faults (Fig. 17). For the same reasons, the fault angle is lower on CDNF margins than on oceanward-dipping normal faults margins. For instance, on the CDNF-controlled margins of Southern Angola and Gabon, we observe that the syn-rift sediments are tilted oceanward by respectively 20° to 35° (Profile ANG1) and 18° to 30° (profile GA2). In contrast, the syn-rift sediments of the Iberian margin, present lower tilt values, circa 5° to 25° (determined after Manatschal, 2004).

#### 4. Conclusion

As a first order key-point we put forward the important variability in the geometry of the present-day rifted margins. Numerous structural parameters such as the length of the margin, the sediment thickness, the dip direction of the normal faults, the abundance of syn-rift material, the inherited thermal state of the lithosphere, and the extension rate during rifting are extremely variable from a margin to another (Fig. 1). To

support this assertion and to investigate further the possible mechanisms responsible for that variability, we have presented a new set of high-resolution seismic reflection profiles from rifted margins of Uruguay, Southern Namibia, Gabon, South China Sea and Barents Sea. This dataset allows us to highlight several tectonic features along these margins:

- Ductile deformation at the base of the crust is imaged on the margins of Uruguay, Gabon and South China Sea. This ductile behavior of a weak lower crust is attributed to an elevated thermal gradient possibly related to (i) a blanketing effect of the syn-rift sediments (Gabon); (ii) to an inherited high thermal gradient (collapsing arc, South China Sea); or (iii) to a plume related thermal anomaly (Uruguay). The frequent weakness of the lower crust, already suggested by earlier numerical models, is now confirmed and clearly imaged by seismic reflection, which should lead to reconsidering models of continental rifting.
- Tectonic boudinage of the continental crust plays an important role during the structuration of the Gabon, Barents, and South China Sea margins. This boudinage is another indication of the weakness of the lower crust. It explains the periodicity of normal faults and the focusing of deformation in the inter-boudin necks. It favors the development of shallow-dipping crustal-scale shear zones and allow for the nucleation of low-angle normal faults. Boudinage and low-angle-normal faults are thus direct consequences of the rheological heterogeneity of the crust and the weakness of the lower crust similar to examples of post-orogenic extension such as the Basin and Range or the Aegean.
- Continentward Dipping Normal Faults (CDNF), probably underestimated until now, are frequent and can control the formation of an entire rifted margin (e.g., Southern Angola, Uruguay...). At the scale of the entire rift (conjugate margins) the fault pattern can be symmetrical, especially in the case of volcanic rifted margins (Fig. 18a). In this case, CDNF are observed on either side, dipping away from the rift centre. Depending on the case, we propose three mechanisms for their formation: i) inversion of inherited faults, that leads to asymmetric conjugate margins; ii) formation above a basal shear possibly related to lithosphere-asthenosphere relative motion, also leading to asymmetric conjugate margins (Fig. 18b); iii) Syn-magmatic gravity collapse above a rising plume head which, in this case, leads to symmetric conjugate volcanic margins (e.g. Uruguay-Namibia margins, Figs. 2 and 18a).

The observation and interpretation of high-quality and deep penetration seismic profiles in several locations leads to an unorthodox vision of rifted margins where the lower crust is weak. Boudinage controls a large part of the deformation and localization of low-angle normal faults, and these normal faults often dip toward the continent (Fig. 18b). Whether these margins are representative of a significant part of rifted margins around the oceans is difficult to assess given the relatively small number of analyzed profiles, but these profiles sample depths that have seldom been imaged before with such details and thus they are the only window we have so far on the behavior of the lower crust in extensional contexts. At the very least, they show that the more classical upper/lower plate asymmetrical models with normal faults dipping toward the ocean, and which involve more resistant rheologies for the lower crust represent merely one category of rifted margins, probably the “cold” ones. “Hot” and hence ductile margins, whether volcanic or not, present a wider distribution of deformation, a lower taper angle, and the characteristics related to a weak lower crust mentioned above (Fig. 18b), while “cold” margins would show a more localized deformation, and a higher taper angle.

#### Acknowledgements

We are grateful to Total SA for funding and offering access to the seismics. We thank Schlumberger Multiclient, Petrolin, TGS, Marathon,

Cobalt and Ion Geophysical for providing outstanding seismic profiles and allowing us to publish them. The Gabonese DGH is thanked for the access to the Gabonese profiles. Emily Clerc greatly helped to improve the English. This work was funded by Total through a postdoc grant to C. Clerc. This paper is a contribution of the ERC Advanced Research Grant RHEOLITH (grant agreement No. 290864), of Institut Universitaire de France and Labex VOLTAIRE.

## References

- Allmendinger, R.W., Hauge, T.A., Hauser, C.J., Potter, S.L., Klempner, K.D., Nelson, P., Knuefer, P., Oliver, J., 1987. Overview of the COCORP 40°N transect, western United States: the fabric of an orogenic belt. *Geological Society of America Bulletin* 98, 308–319.
- Aslanian, D., Moulin, M., 2013. Palaeogeographic consequences of conservative models in the South Atlantic Ocean. *Geol. Soc. Lond. Spec. Publ.* 369:75–90. <http://dx.doi.org/10.1144/SP369.5>.
- Aslanian, D., Moulin, M., Olivet, J.-L., Unternehr, P., Matias, L., Bache, F., Rabineau, M., Nouzé, H., Klingelhoefer, F., Contrucci, I., Labails, C., 2009. Brazilian and African passive margins of the Central Segment of the South Atlantic Ocean: kinematic constraints. *Tectonophysics* 468:98–112. Role of Magmatism in Continental Lithosphere Extension. [10.1016/j.tecto.2008.12.016](http://dx.doi.org/10.1016/j.tecto.2008.12.016).
- Autin, J., Leroy, S., Beslier, M.-O., D'Acremont, E., Razin, P., Ribodetti, A., Bellahsen, N., Robin, C., Al Toubi, K., 2010. Continental break-up history of a deep magma-poor margin based on seismic reflection data (northeastern Gulf of Aden margin, offshore Oman). *Geophysical Journal International* 180:501–519. <http://dx.doi.org/10.1111/j.1365-246X.2009.04424.x>.
- Barckhausen, U., Roeser, H.A., 2004. Seafloor spreading anomalies in the South China Sea revisited. In: Clift, P., Kuhnt, W., Wang, P., Hayes, D. (Eds.), *Continental-Ocean Interactions Within East Asian Marginal Seas*. American Geophysical Union, pp. 121–125.
- Barton, A.J., White, R.S., 1997. Volcanism on the Rockall continental margin. *Journal of the Geological Society* 154:531–536. <http://dx.doi.org/10.1144/gsjgs.154.3.0531>.
- Becker, K., Tanner, D.C., Franke, D., Krawczyk, C.M., 2016. Fault-controlled lithospheric detachment of the volcanic southern South Atlantic rift. *Geochemistry, Geophysics, Geosystems* <http://dx.doi.org/10.1002/2015GC006081> (n/a–n/a).
- Bialas, R.W., Buck, W.R., 2009. How sediment promotes narrow rifting: application to the Gulf of California. *Tectonics* 28, TC4014. <http://dx.doi.org/10.1029/2008TC002394>.
- Blaich, O.A., Faleide, J.I., Tsikalas, F., Gordon, A.C., Mohriak, W., 2013. Crustal-scale architecture and segmentation of the South Atlantic volcanic margin. *Geological Society of London, Special Publication* 369:167–183. <http://dx.doi.org/10.1144/SP369.22>.
- Boillot, G., Winterer, E.L., Meyer, A.W., et al. (Eds.), 1987. *Proceedings of the Ocean Drilling Program, 103 Initial Reports*. Ocean Drilling Program.
- Breivik, A.J., Faleide, J.I., Gudlaugsson, S.T., 1998. Southwestern Barents Sea margin: late Mesozoic sedimentary basins and crustal extension. *Tectonophysics* 293:21–44. [http://dx.doi.org/10.1016/S0040-1951\(98\)00073-0](http://dx.doi.org/10.1016/S0040-1951(98)00073-0).
- Briaes, A., Patriat, P., Tapponnier, P., 1993. Updated interpretation of magnetic anomalies and seafloor spreading stages in the south China Sea: implications for the Tertiary tectonics of Southeast Asia. *Journal of Geophysical Research - Solid Earth* 98: 6299–6328. <http://dx.doi.org/10.1029/92JB02280>.
- Brito, M., Figueiredo, C., Inkkollu, M., 2004. Namibe Basin: deepwater exploration potential. *AAPG Int. Conf. Exhib.* 14–18 Novemb. 2004 Abuja Niger.
- Brooks, C.K., Nielsen, T.F.D., 1982. The E Greenland continental margin: a transition between oceanic and continental magmatism. *Journal of the Geological Society* 139: 265–275. <http://dx.doi.org/10.1144/gsjgs.139.3.0265>.
- Brun, J.P., Beslier, M.O., 1996. Mantle exhumation at passive margins. *Earth and Planetary Science Letters* 142:161–173. [http://dx.doi.org/10.1016/0012-821X\(96\)00080-5](http://dx.doi.org/10.1016/0012-821X(96)00080-5).
- Brun, J.-P., Choukroune, P., Faugeres, E., 1985. Les discontinuités significatives de l'amincissement crustal; application aux marges passives. *Bulletin de la Société Géologique de France* 1:139–144. <http://dx.doi.org/10.2113/gssgfbull.1.1.139>.
- Brune, S., Heine, C., Pérez-Gussinyé, M., Sobolev, S.V., 2014. Rift migration explains continental margin asymmetry and crustal hyper-extension. *Nature Communications* 5. <http://dx.doi.org/10.1038/ncomms5014>.
- Burchfiel, B.C., Hodges, K.V., Royden, L.H., 1987. *Geology of Panamint Valley-Saline Valley pull-apart system, California; palinspastic evidence for low-angle geometry of a Neogene range-bounding fault*. *Journal of Geophysical Research* 92, 10422–10426.
- Burrus, J., 1984. Contribution to a geodynamic synthesis of the Provençal Basin (North-Western Mediterranean). *Marine Geology* 55:247–269. [http://dx.doi.org/10.1016/0025-3227\(84\)90071-9](http://dx.doi.org/10.1016/0025-3227(84)90071-9).
- Chen, Y., Niu, F., Liu, R., Huang, Z., Kralčić, H., Sun, L., Chan, W., 2010. Crustal structure beneath China from receiver function analysis. *Journal of Geophysical Research - Solid Earth* 115, B03307. <http://dx.doi.org/10.1029/2009JB006386>.
- Clerc, C., Lagabrielle, Y., 2014. Thermal control on the modes of crustal thinning leading to mantle exhumation. Insights from the Cretaceous Pyrenean hot paleomargins. *Tectonics* <http://dx.doi.org/10.1002/2013TC003471> (2013TC003471).
- Clerc, C., Jolivet, L., Ringenbach, J.-C., 2015a. Ductile extensional shear zones in the lower crust of a passive margin. *Earth and Planetary Science Letters* 431:1–7. <http://dx.doi.org/10.1016/j.epsl.2015.08.038>.
- Clerc, C., Lahfid, A., Monié, P., Lagabrielle, Y., Chopin, C., Poujol, M., Boulvais, P., Ringenbach, J.-C., Masini, E., de St Blanquat, M., 2015b. High-temperature metamorphism during extreme thinning of the continental crust: a reappraisal of the North Pyrenean passive paleomargin. *Solid Earth* 6:643–668. <http://dx.doi.org/10.5194/se-6-643-2015>.
- Clerc, C., Lagabrielle, Y., Labaume, P., Ringenbach, J.-C., Vauchez, A., Nalpas, T., Bousquet, R., Ballard, J.-F., Lahfid, A., Fourcade, S., 2016. Basement – cover decoupling and progressive exhumation of metamorphic sediments at hot rifted margin. Insights from the North-eastern Pyrenean analog. *Tectonophysics* <http://dx.doi.org/10.1016/j.tecto.2016.07.022>.
- Colletini, C., 2011. The mechanical paradox of low-angle normal faults: current understanding and open questions. *Tectonophysics* 510:253–268. <http://dx.doi.org/10.1016/j.tecto.2011.07.015>.
- Contrucci, I., Matias, L., Moulin, M., Géli, L., Klingelhoefer, F., Nouzé, H., Aslanian, D., Olivet, J.L., Réhault, J.-P., Sibuet, J.-C., 2004. Deep structure of the West African continental margin (Congo, Zaïre, Angola), between 5°S and 8°S, from reflection/refraction seismics and gravity data. *Geophysical Journal International* 158:529–553. <http://dx.doi.org/10.1111/j.1365-246X.2004.02303.x>.
- Corre, B., Lagabrielle, Y., Labaume, P., Fourcade, S., Clerc, C., Ballèvre, M., 2016. Deformation associated with mantle exhumation in a distal, hot passive margin environment: new constraints from the Sarailé Massif (Chaînons Béarnais, North-Pyrenean Zone). *Comptes Rendus Geoscience* 348:279–289. <http://dx.doi.org/10.1016/j.crte.2015.11.007>.
- Courtilot, V., 1982. Propagating rifts and continental breakup. *Tectonics* 1:239–250. <http://dx.doi.org/10.1029/TC001i003p00239>.
- Crosby, A., White, N., Edwards, G., Shillington, D.J., 2008. Evolution of the Newfoundland-Iberia conjugate rifted margins. *Earth and Planetary Science Letters* 273:214–226. <http://dx.doi.org/10.1016/j.epsl.2008.06.039>.
- Cullen, A., Reemst, P., Henstra, G., Gozzard, S., Ray, A., 2010. Rifting of the South China Sea: new perspectives. *Petroleum Geoscience* 16:273–282. <http://dx.doi.org/10.1144/1354-079309-908>.
- Davis, M., Kusznir, N., 2004. Depth-dependent lithospheric stretching at rifted continental margins. In: Karner, G.D. (Ed.), *Proceedings of NSF Rifted Margins Theoretical Institute*, pp. 92–137 United States.
- Ding, W., Li, J., 2016. Propagated rifting in the Southwest Sub-basin, South China Sea: insights from analogue modelling. *Journal of Geodynamics* <http://dx.doi.org/10.1016/j.jog.2016.02.004>.
- Driscoll, N.W., Karner, G.D., 1998. Lower crustal extension across the Northern Carnarvon basin, Australia: evidence for an eastward dipping detachment. *Journal of Geophysical Research - Solid Earth* 103:4975–4991. <http://dx.doi.org/10.1029/97JB03295>.
- Dupré, S., Bertotti, G., Cloetingh, S., 2007. Tectonic history along the South Gabon Basin: anomalous early post-rift subsidence. *Marine and Petroleum Geology* 24:151–172. <http://dx.doi.org/10.1016/j.marpetgeo.2006.11.003>.
- Dupré, S., Cloetingh, S., Bertotti, G., 2011. Structure of the Gabon Margin from integrated seismic reflection and gravity data. *Tectonophysics* 506:31–45. <http://dx.doi.org/10.1016/j.tecto.2011.04.009>.
- Egan, S.S., Meredith, D.J., 2007. A kinematic modelling approach to lithosphere deformation and basin formation: application to the Black Sea. *Geological Society of London, Special Publication* 282:173–198. <http://dx.doi.org/10.1144/SP282.9>.
- Faleide, J.I., Gudlaugsson, S.T., Jacquart, G., 1984. Evolution of the western Barents Sea. *Marine and Petroleum Geology* 1:123–150. [http://dx.doi.org/10.1016/0264-8172\(84\)90082-5](http://dx.doi.org/10.1016/0264-8172(84)90082-5).
- Faleide, J.I., Vågenes, E., Gudlaugsson, S.T., 1993. Late Mesozoic-Cenozoic evolution of the south-western Barents Sea in a regional rift-shear tectonic setting. *Marine and Petroleum Geology* 10:186–214. [http://dx.doi.org/10.1016/0264-8172\(93\)90104-Z](http://dx.doi.org/10.1016/0264-8172(93)90104-Z).
- Famin, V., Philippot, P., Jolivet, L., Agard, P., 2004. Evolution of hydrothermal regime along a crustal shear zone, Tinos island, Greece. *Tectonics* 23. <http://dx.doi.org/10.1029/2003TC001509>.
- Faugeres, E., 1985. *La Tectonique en extension intracontinentale: étude de terrain, le Sud du Nevada, U.S.A., et modélisation analogique*.
- Faugeres, E., Brun, J.-P., 1984. Modélisation expérimentale de la distension continentale. *Comptes Rendus. Académie des Sciences* 299, 365.
- Fletcher, R.C., Hallet, B., 1983. Unstable extension of the lithosphere: a mechanical model for basin-and-range structure. *Journal of Geophysical Research* 88, 7457–7466.
- Franke, D., Neben, S., Schreckenberger, B., Schulze, A., Stiller, M., Krawczyk, C.M., 2006. Crustal structure across the Colorado Basin, offshore Argentina. *Geophysical Journal International* 165:850–864. <http://dx.doi.org/10.1111/j.1365-246X.2006.02907.x>.
- Franke, D., Neben, S., Ladage, S., Schreckenberger, B., Hinz, K., 2007. Margin segmentation and volcano-tectonic architecture along the volcanic margin off Argentina/Uruguay, South Atlantic. *Marine Geology* 244:46–67. <http://dx.doi.org/10.1016/j.margeo.2007.06.009>.
- Franke, D., Savva, D., Pubellier, M., Steuer, S., Mouly, B., Auxietre, J.-L., Meresse, F., Chamot-Rooke, N., 2014. The final rifting evolution in the South China Sea. *Marine and Petroleum Geology* 58 (Part B):704–720. Evolution, Structure, and Sedimentary Record of the South China Sea and Adjacent Basins. [10.1016/j.marpetgeo.2013.11.020](http://dx.doi.org/10.1016/j.marpetgeo.2013.11.020).
- Froidevaux, C., 1986. Basin and range large-scale tectonics: constraints from gravity and reflection seismology. *Journal of Geophysical Research* 91, 3625–3632.
- Fromm, T., Planert, L., Jokat, W., Ryberg, T., Behrmann, J.H., Weber, M.H., Haberland, C., 2015. South Atlantic opening: a plume-induced breakup? *Geology* 43:931–934. <http://dx.doi.org/10.1130/G36936.1>.
- Gartrell, A.P., 1997. Evolution of rift basins and low-angle detachments in multilayer analog models. *Geology* 25:615–618. [http://dx.doi.org/10.1130/0091-7613\(1997\)025<0615: EORBAL>2.3.CO;2](http://dx.doi.org/10.1130/0091-7613(1997)025<0615: EORBAL>2.3.CO;2).
- Geoffroy, L., 2001. The structure of volcanic margins: some problematics from the North-Atlantic/Labrador-Baffin system. *Marine and Petroleum Geology* 18: 463–469. [http://dx.doi.org/10.1016/S0264-8172\(00\)00073-8](http://dx.doi.org/10.1016/S0264-8172(00)00073-8).
- Geoffroy, L., 2005. Volcanic passive margins. *Comptes Rendus Geoscience* 337: 1395–1408. <http://dx.doi.org/10.1016/j.crte.2005.10.006>.
- Geoffroy, L., Callot, J.-P., Scaillet, S., Skuce, A., Gélard, J.P., Ravilly, M., Angelier, J., Bonin, B., Cayet, C., Perrot, K., Lepvrier, C., 2001. Southeast Baffin volcanic margin and the North American-Greenland plate separation. *Tectonics* 20:566–584. <http://dx.doi.org/10.1029/2001TC900003>.
- Geoffroy, L., Burov, E.B., Werner, P., 2015. Volcanic passive margins: another way to break up continents. *Scientific Reports* 5. <http://dx.doi.org/10.1038/srep14828>.

- Gernigon, L., Ringenbach, J.-C., Planke, S., Le Gall, B., 2004. Deep structures and breakup along volcanic rifted margins: insights from integrated studies along the outer Vøring Basin (Norway). *Marine and Petroleum Geology* 21:363–372. <http://dx.doi.org/10.1016/j.marpetgeo.2004.01.005>.
- Gernigon, L., Lucazeau, F., Brigaud, F., Ringenbach, J.-C., Planke, S., Le Gall, B., 2006. A moderate melting model for the Vøring margin (Norway) based on structural observations and a thermo-kinematic modelling: implication for the meaning of the lower crustal bodies. *Tectonophysics* 412:255–278. <http://dx.doi.org/10.1016/j.tecto.2005.10.038>.
- Gernigon, L., Brønner, M., Roberts, D., Olesen, O., Nasuti, A., Yamasaki, T., 2014. Crustal and basin evolution of the southwestern Barents Sea: from Caledonian orogeny to continental breakup. *Tectonics* 33. <http://dx.doi.org/10.1002/2013TC003439> (2013TC003439).
- Gibson, S.A., Thompson, R.N., Day, J.A., 2006. Timescales and mechanisms of plume–lithosphere interactions: 40Ar/39Ar geochronology and geochemistry of alkaline igneous rocks from the Paraná–Etendeka large igneous province. *Earth and Planetary Science Letters* 251:1–17. <http://dx.doi.org/10.1016/j.epsl.2006.08.004>.
- Gładczenko, T.P., Hinz, K., Eldholm, O., Meyer, H., Neben, S., Skogseid, J., 1997. South Atlantic volcanic margins. *Journal of the Geological Society* 154:465–470. <http://dx.doi.org/10.1144/gsjgs.154.3.0465>.
- Gładczenko, T.P., Skogseid, J., Eldholm, O., 1998. Namibia volcanic margin. *Marine Geophysical Researches* 20:313–341. <http://dx.doi.org/10.1023/A:1004746101320>.
- Goetze, C., Evans, B., 1979. Stress and temperature in the bending lithosphere as constrained by experimental rock mechanics. *Geophysical Journal of the Royal Astronomical Society* 59, 463–478.
- Goscombe, B.D., Passchier, C.W., Hand, M., 2004. Boudinage classification: end-member boudin types and modified boudin structures. *Journal of Structural Geology* 26, 739–763.
- Griffiths, R.W., Campbell, I.H., 1991. Interaction of mantle plume heads with the Earth's surface and onset of small-scale convection. *Journal of Geophysical Research - Solid Earth* 96:18295–18310. <http://dx.doi.org/10.1029/91JB01897>.
- Gudlaugsson, S.T., Faleide, J.J., Johansen, S.E., Breivik, A.J., 1998. Late Palaeozoic structural development of the South-western Barents Sea. *Marine and Petroleum Geology* 15: 73–102. [http://dx.doi.org/10.1016/S0264-8172\(97\)00048-2](http://dx.doi.org/10.1016/S0264-8172(97)00048-2).
- Handy, M.R., Zingg, A., 1991. The tectonic and rheological evolution of an attenuated cross section of the continental crust: Ivrea crustal section, southern Alps, northwestern Italy and southern Switzerland. *Geological Society of America Bulletin* 103: 236–253. [http://dx.doi.org/10.1130/0016-7606\(1991\)103<0236:TTAREO>2.3.CO;2](http://dx.doi.org/10.1130/0016-7606(1991)103<0236:TTAREO>2.3.CO;2).
- Harry, D.L., Sawyer, D.S., 1992. A dynamic model of extension in the Baltimore Canyon Trough Region. *Tectonics* 11:420–436. <http://dx.doi.org/10.1029/91TC03012>.
- Hauptert, I., Manatschal, G., Decarlis, A., Unternehr, P., 2016. Upper-plate magma-poor rifted margins: stratigraphic architecture and structural evolution. *Marine and Petroleum Geology* 69:241–261. <http://dx.doi.org/10.1016/j.marpetgeo.2015.10.020>.
- Heine, C., Zoethout, J., Müller, R.D., 2013. Kinematic of the South Atlantic rift. *Solid Earth* 4: 215–253. <http://dx.doi.org/10.5194/se-4-215-2013>.
- Hinz, K., 1981. A Hypothesis on Terrestrial Catastrophes: Wedges of Very Thick Oceanward Dipping Layers Beneath Passive Continental Margins; Their Origin and Paleoenvironmental Significance. *Schweizerbart*.
- Hirsch, K.K., Bauer, K., Scheck-Wenderoth, M., 2009. Deep structure of the western South African passive margin – results of a combined approach of seismic, gravity and isostatic investigations. *Tectonophysics* 470:57–70. *Progress in Understanding Sedimentary Basins*. 10.1016/j.tecto.2008.04.028.
- Hopper, J.R., Buck, W.R., 1996. The effect of lower crustal flow on continental extension and passive margin formation. *Journal of Geophysical Research - Solid Earth* 101: 20175–20194. <http://dx.doi.org/10.1029/96JB01644>.
- Huchon, P., Nguyen, T.N.H., Chamot-Rooke, N., 2001. Propagation of continental break-up in the southwestern South China Sea. *Geological Society of London, Special Publication* 187:31–50. <http://dx.doi.org/10.1144/GSL.SP.2001.187.01.03>.
- Huismans, R., Beaumont, C., 2011. Depth-dependent extension, two-stage breakup and cratonic underplating at rifted margins. *Nature* 473:74–78. <http://dx.doi.org/10.1038/nature09988>.
- Huismans, R.S., Beaumont, C., 2014. Rifted continental margins: the case for depth-dependent extension. *Earth and Planetary Science Letters* 407:148–162. <http://dx.doi.org/10.1016/j.epsl.2014.09.032>.
- Jackson, M.P.A., Cramez, C., Fonck, J.-M., 2000. Role of subaerial volcanic rocks and mantle plumes in creation of South Atlantic margins: implications for salt tectonics and source rocks. *Marine and Petroleum Geology* 17:477–498. [http://dx.doi.org/10.1016/S0264-8172\(00\)00066-4](http://dx.doi.org/10.1016/S0264-8172(00)00066-4).
- Jammes, S., Lavier, L.L., 2016. The effect of bimineralic composition on extensional processes at lithospheric scale. *Geochemistry, Geophysics, Geosystems* 17:3375–3392. <http://dx.doi.org/10.1002/2016GC006399>.
- Jolivet, L., Famin, V., Mehl, C., Parra, T., Aubourg, C., Hébert, R., Philippot, P., 2004. Strain localization during crustal-scale boudinage to form extensional metamorphic domes in the Aegean Sea. *Geological Society of America Special Papers* 380: 185–210. <http://dx.doi.org/10.1130/G01837-2380-9.185>.
- Jolivet, L., Labrousse, L., Agard, P., Lacombe, O., Bailly, V., Lecomte, E., Mouthereau, F., Mehl, C., 2010. Rifting and shallow-dipping detachments, clues from the Corinth Rift and the Aegean. *Tectonophysics* 483:287–304. <http://dx.doi.org/10.1016/j.tecto.2009.11.001>.
- Jolivet, L., Gorini, C., Smit, J., Leroy, S., 2015. Continental breakup and the dynamics of rifting in back-arc basins: the Gulf of Lion margin. *Tectonics* 34. <http://dx.doi.org/10.1002/2014TC003570> (2014TC003570).
- Karner, G.D., 1991. Sediment blanketing and the flexural strength of extended continental lithosphere. *Basin Research* 3:177–185. <http://dx.doi.org/10.1111/j.1365-2117.1991.tb00127.x>.
- Karner, G.D., Gambôa, L.A.P., 2007. Timing and origin of the South Atlantic pre-salt sag basins and their capping evaporites. *Geological Society of London, Special Publication* 285:15–35. <http://dx.doi.org/10.1144/SP285.2>.
- Lacombe, O., Jolivet, L., Le Pourhiet, L., Lecomte, E., Mehl, C., 2013. Initiation, geometry and mechanics of brittle faulting in exhuming metamorphic rocks: insights from the northern Cycladic islands (Aegean, Greece). *Bulletin de la Société Géologique de France* 184, 383–403.
- Lagabrielle, Y., Clerc, C., Vauchez, A., Lahfid, A., Labaume, P., Azambre, B., Fourcade, S., Dautria, J.-M., 2016. Very high geothermal gradient during mantle exhumation recorded in mylonitic marbles and carbonate breccias from a Mesozoic Pyrenean palaeomargin (Lherz area, North Pyrenean Zone, France). *Comptes Rendus Geoscience* 348:290–300. <http://dx.doi.org/10.1016/j.crte.2015.11.004>.
- Lavier, L.L., Manatschal, G., 2006. A mechanism to thin the continental lithosphere at magma-poor margins. *Nature* 440:324–328. <http://dx.doi.org/10.1038/nature04608>.
- Lavier, L.L., Steckler, M.S., 1997. The effect of sedimentary cover on the flexural strength of continental lithosphere. *Nature* 389:476–479. <http://dx.doi.org/10.1038/39004>.
- Le Pichon, X., Chamot-Rooke, N., 1991. Extension of Continental Crust, *Controversies in Modern Geology*. Academic Press Limited, New-York, pp. 313–338.
- Lecomte, E., Le Pourhiet, L., Lacombe, O., Jolivet, L., 2011. A continuum mechanics approach to quantify brittle strain on weak faults: application to the extensional reactivation of shallow-dipping discontinuities. *Geophysical Journal International* 184: 1–11. <http://dx.doi.org/10.1111/j.1365-1246X.2010.04821.x>.
- Lentini, M.R., Fraser, S.L., Sumner, H.S., Davies, R.J., 2010. Geodynamics of the central South Atlantic conjugate margins: implications for hydrocarbon potential. *Petroleum Geoscience* 16:217–229. <http://dx.doi.org/10.1144/1354-079309-909>.
- Levell, B., Argent, J., Doré, A.G., Fraser, S., 2010. Passive margins: overview. *Geological Society of London* 7:823–830. *Pet. Geol. Conf. Ser.* 10.1144/0070823.
- Lister, G.S., Etheridge, M.A., Symonds, P.A., 1986. Detachment faulting and the evolution of passive continental margins. *Geology* 14:246–250. [http://dx.doi.org/10.1130/0091-7613\(1986\)14<246:DFATEO>2.0.CO;2](http://dx.doi.org/10.1130/0091-7613(1986)14<246:DFATEO>2.0.CO;2).
- Lizarralde, D., Axen, G.J., Brown, H.E., Fletcher, J.M., González-Fernández, A., Harding, A.J., Holbrook, W.S., Kent, G.M., Paramo, P., Sutherland, F., Umhoefer, P.J., 2007. Variation in styles of rifting in the Gulf of California. *Nature* 448:466–469. <http://dx.doi.org/10.1038/nature06035>.
- Lohest, M., 1909. L'origine des veines et des géodes des terrains primaires de Belgique. *Annales de la Société Géologique de Belgique* 36B, 275–282.
- Magnavita, L.P., Davison, I., Kuszniir, N.J., 1994. Rifting, erosion, and uplift history of the Recôncavo-Tucano-Jatobá Rift, northeast Brazil. *Tectonics* 13:367–388. <http://dx.doi.org/10.1029/93TC02941>.
- Manatschal, G., 2004. New models for evolution of magma-poor rifted margins based on a review of data and concepts from West Iberia and the Alps. *International Journal of Earth Sciences* 93:432–466. <http://dx.doi.org/10.1007/s00531-004-0394-7>.
- Marrett, R., Allmendinger, R.W., 1992. Amount of extension on "small" faults: an example from the Viking graben. *Geology* 20:47–50. [http://dx.doi.org/10.1130/0091-7613\(1992\)020<0047:AOE0SF>2.3.CO;2](http://dx.doi.org/10.1130/0091-7613(1992)020<0047:AOE0SF>2.3.CO;2).
- Martin, J., Toothill, S., Veritas, C.G.G., Moussavou, R., 2009. *Hunting the pre-salt*. *GeoExpro* 6, 36–41.
- Maus, S., Barckhausen, U., Berkenbosch, H., Bournas, N., Brozena, J., Childers, V., Dostaler, F., Fairhead, J.D., Finn, C., von Frese, R.R.B., Gaina, C., Golynsky, S., Kucks, R., Lühr, H., Milligan, P., Mogren, S., Müller, R.D., Olesen, O., Pilkington, M., Saltus, R., Schreckenberger, B., Thébaud, E., Caratori Tontini, F., 2009. EMAG2: a 2-arc min resolution Earth Magnetic Anomaly Grid compiled from satellite, airborne, and marine magnetic measurements. *Geochemistry, Geophysics, Geosystems* 10, Q08005. <http://dx.doi.org/10.1029/2009GC002471>.
- Mbina Mounquengui, M., Guiraud, M., 2009. Neocomian to early Aptian syn-rift evolution of the normal to oblique-rifted North Gabon Margin (Interior and N'Komi Basins). *Marine and Petroleum Geology* 26:1000–1017. <http://dx.doi.org/10.1016/j.marpetgeo.2008.11.001>.
- McDermott, K., Gillard, E., Clarke, N., 2015. From basalt to skeletons – the 200 million-year history of the Namibian margin uncovered by new seismic data. *First Break* 33, 77–85.
- Mehl, C., Jolivet, L., Lacombe, O., 2005. From ductile to brittle: evolution and localization of deformation below a crustal detachment (Tinos, Cyclades, Greece). *Tectonics* 24 (doi: 4010.1029/2004TC001767. TC4017).
- Mehl, C., Jolivet, L., Lacombe, O., Labrousse, L., Rimmelé, G., 2007. Structural evolution of Andros (Cyclades, Greece): a key to the behaviour of a (flat) detachment within an extending continental crust. *Geological Society of London, Special Publication* 291: 41–73. <http://dx.doi.org/10.1144/SP291.3>.
- Mello, M.R., Filho, N.C.D.A., Bender, A.A., Barbanti, S.M., Mohriak, W., Schmitt, P., Jesus, C.L.C.D., 2013. The Namibian and Brazilian southern South Atlantic petroleum systems: are they comparable analogues? *Geological Society of London, Special Publication* 369:249–266. <http://dx.doi.org/10.1144/SP369.18>.
- Meyers, J.B., Rosendahl Jr., B.R., Austin, J.A., 1996. Deep-penetrating MCS images of the South Gabon Basin: implications for rift tectonics and post-breakup salt remobilization. *Basin Research* 8:65–84. <http://dx.doi.org/10.1111/j.1365-2117.1996.tb00115.x>.
- Michon, L., Merle, O., 2003. Mode of lithospheric extension: conceptual models from analogue modeling. *Tectonics* 22:1028. <http://dx.doi.org/10.1029/2002TC001435>.
- Mjelde, R., Raum, T., Murai, Y., Takanami, T., 2007. Continent-ocean-transitions: review, and a new tectono-magmatic model of the Vøring Plateau, NE Atlantic. *Journal of Geodynamics* 43:374–392. <http://dx.doi.org/10.1016/j.jog.2006.09.013>.
- Mohn, G., Manatschal, G., Beltrando, M., Masini, E., Kuszniir, N., 2012. Necking of continental crust in magma-poor rifted margins: evidence from the fossil Alpine Tethys margins: necking of continental crust. *Tectonics* 31. <http://dx.doi.org/10.1029/2011TC002961> (n/a-n/a).
- Mohriak, W.U., Lira Rabelo, J., De Matos, R.D., De Barros, M.C., 1995. Deep seismic reflection profiling of sedimentary basins offshore Brazil: geological objectives and preliminary results in the Sergipe Basin. *Journal of Geodynamics* 20:515–539. *Proceedings of the International Symposium on the Physics and Chemistry of the Upper Mantle*. 10.1016/0264-3707(95)00024-4.
- Mohriak, W.U., Nemcok, M., Enciso, G., 2008. South Atlantic divergent margin evolution: rift-border uplift and salt tectonics in the basins of SE Brazil. In: Pankhurst, R., Trouw, R.A., Brito Neves, B.B., de Wit, M. (Eds.), *West Gondwana Pre-Cenozoic Correlations Across*

- the South Atlantic Region. Geological Society, London, Special Publications 294, pp. 365–398.
- Mohriak, W.U., Mello, M.R., and Azambuja Filho, N.C., 2012. Tectonic reconstructions and petroleum system correlations in the southern south Atlantic: exploratory analysis and the search for a new hydrocarbon play. In: N.C. Rosen et al. (eds.), *New Understanding of the Petroleum Systems of Continental Margins of the World*, 32nd Annual GCSS-EPM Foundation Bob F. Perkins Research Conference, December 2–5, 2012, Houston, Texas, Expanded Abstracts, 306–346.
- Moulin, M., Aslanian, D., Olivet, J.-L., Contrucci, I., Matias, L., Géli, L., Klingelhoefer, F., Nouzé, H., Réhault, J.-P., Unternehr, P., 2005. Geological constraints on the evolution of the Angolan margin based on reflection and refraction seismic data (ZaiAngo project). *Geophysical Journal International* 162:793–810. <http://dx.doi.org/10.1111/j.1365-246X.2005.02668.x>.
- Moulin, M., Aslanian, D., Rabineau, M., Patriat, M., Matias, L., 2013. Kinematic keys of the Santos-Namibe basins. Geological Society of London, Special Publication 369:91–107. <http://dx.doi.org/10.1144/SP369.3>.
- Müntener, O., Hermann, J., 2001. The role of lower crust and continental upper mantle during formation of non-volcanic passive margins: evidence from the Alps. Geological Society of London, Special Publication 187:267–288. <http://dx.doi.org/10.1144/GSLSP.2001.187.01.13>.
- Mutter, J.C., 1985. Geophysics of the Polar Regions Seaward dipping reflectors and the continent-ocean boundary at passive continental margins. *Tectonophysics* 114: 117–131. [http://dx.doi.org/10.1016/0040-1951\(85\)90009-5](http://dx.doi.org/10.1016/0040-1951(85)90009-5).
- Mutter, J.C., Talwani, M., Stoffa, P.L., 1982. Origin of seaward-dipping reflectors in oceanic crust off the Norwegian margin by "subaerial sea-floor spreading". *Geology* 10:353. [http://dx.doi.org/10.1130/0091-7613\(1982\)10<353:OOSRIO>2.0.CO;2](http://dx.doi.org/10.1130/0091-7613(1982)10<353:OOSRIO>2.0.CO;2).
- Nissen, S.S., Hayes, D.E., Bochu, Y., Weijun, Zeng, Yongqin, Chen, Xiaupin, Nu, 1995. Gravity, heat flow, and seismic constraints on the processes of crustal extension: northern margin of the South China Sea. *Journal of Geophysical Research* 100, 22447–22483.
- Osmundsen, P.T., Andersen, T.B., 2001. The middle Devonian basins of western Norway: sedimentary response to large-scale transtensional tectonics? *Tectonophysics* 332, 51–68.
- Osmundsen, P.T., Ebbing, J., 2008. Styles of extension offshore mid-Norway and implications for mechanisms of crustal thinning at passive margins. *Tectonics* 27, TC6016. <http://dx.doi.org/10.1029/2007TC002242>.
- Pérez-Gussinyé, M., Ranero, C.R., Reston, T.J., Sawyer, D., 2003. Mechanisms of extension at nonvolcanic margins: evidence from the Galicia interior basin, west of Iberia. *Journal of Geophysical Research - Solid Earth* 108. <http://dx.doi.org/10.1029/2001JB000901> (n/a–n/a).
- Péron-Pinvidic, G., Manatschal, G., 2008. The final rifting evolution at deep magma-poor passive margins from Iberia-Newfoundland: a new point of view. *International Journal of Earth Sciences* 98:1581–1597. <http://dx.doi.org/10.1007/s00531-008-0337-9>.
- Péron-Pinvidic, G., Manatschal, G., Minshull, T.A., Sawyer, D.S., 2007. Tectono-sedimentary evolution of the deep Iberia-Newfoundland margins: evidence for a complex breakup history. *Tectonics* 26, TC2011. <http://dx.doi.org/10.1029/2006TC001970>.
- Péron-Pinvidic, G., Manatschal, G., Masini, E., Sutra, E., Flament, J.M., Hauptert, I., Unternehr, P., 2015. Unravelling the along-strike variability of the Angola–Gabon rifted margin: a mapping approach. Geological Society of London, Special Publication 438. <http://dx.doi.org/10.1144/SP438.1> (SP438.1).
- Pichot, T., Delescluse, M., Chamot-Rooke, N., Pubellier, M., Qiu, Y., Meresse, F., Sun, G., Savva, D., Wong, K.P., Watremez, L., Auxietre, J.L., 2014. Deep crustal structure of the conjugate margins of the SW South China Sea from wide-angle refraction seismic data. *Marine and Petroleum Geology* 58, 627–643.
- Pindell, J., Graham, R., Horn, B., 2014. Rapid outer marginal collapse at the rift to drift transition of passive margin evolution, with a Gulf of Mexico case study. *Basin Research* 26:701–725. <http://dx.doi.org/10.1111/bre.12059>.
- Planke, S., Symonds, P.A., Alvestad, E., Skogseid, J., 2000. Seismic volcanostratigraphy of large-volume basaltic extrusive complexes on rifted margins. *Journal of Geophysical Research - Solid Earth* 105:19335–19351. <http://dx.doi.org/10.1029/1999JB900005>.
- Proffett, J.M., 1977. Cenozoic geology of the Yerington district, Nevada, and implications for the nature and origin of basin and range faulting. Geological Society of America Bulletin 88:247–266. [http://dx.doi.org/10.1130/0016-7606\(1977\)88<247:CGOTYD>2.0.CO;2](http://dx.doi.org/10.1130/0016-7606(1977)88<247:CGOTYD>2.0.CO;2).
- Pubellier, M., Ego, F., Chamot-Rooke, N., Rangin, C., 2003. The building of pericratonic mountain ranges: structural and kinematic constraints applied to GIS-based reconstructions of SE Asia. *Bulletin de la Societe Geologique de France* 174:561–584. <http://dx.doi.org/10.2113/174.6.561>.
- Rabinowitz, P.D., LaBrecque, J.L., 1979. The Mesozoic South Atlantic Ocean and evolution of its continental margins. *Journal of Geophysical Research* 84:5973–6002. <http://dx.doi.org/10.1029/JB084IB11p05973>.
- Raillard, S., Biteau, J.J., All, P., 1998. Abstract: Lower Congo Tertiary Basin - offshore West Africa structural zonation and evolution. AAPG Bulletin 82 (1998). <http://dx.doi.org/10.1306/1D9BD7B3-172D-11D7-8645000102C1865D>.
- Ranero, C.R., Pérez-Gussinyé, M., 2010. Sequential faulting explains the asymmetry and extension discrepancy of conjugate margins. *Nature* 468:294–299. <http://dx.doi.org/10.1038/nature09520>.
- Reston, T.J., 1988. Evidence for shear zones in the lower crust offshore Britain. *Tectonics* 7: 929–945. <http://dx.doi.org/10.1029/TC007i05p0929>.
- Reston, T.J., 2005. Polyphase faulting during the development of the west Galicia rifted margin. *Earth and Planetary Science Letters* 237:561–576. <http://dx.doi.org/10.1016/j.epsl.2005.06.019>.
- Reston, T.J., 2007. The formation of non-volcanic rifted margins by the progressive extension of the lithosphere: the example of the West Iberian margin. Geological Society of London, Special Publication 282:77–110. <http://dx.doi.org/10.1144/SP282.5>.
- Reston, T.J., McDermott, K.G., 2011. Successive detachment faults and mantle unroofing at magma-poor rifted margins. *Geology* 39:1071–1074. <http://dx.doi.org/10.1130/G32428.1>.
- Reston, T.J., McDermott, K., 2014. An assessment of the cause of the "extension discrepancy" with reference to the west Galicia margin. *Basin Research* 26:135–153. <http://dx.doi.org/10.1111/bre.12042>.
- Ricard, Y., Froidevaux, C., 1986. Stretching instabilities and lithospheric boudinage. *Journal of Geophysical Research* 91, 8314–8324.
- Ricard, Y., Froidevaux, C., Simpson, R., 1987. Spectral analysis of topography and gravity in the Basin and Range Province. *Tectonophysics* 133:175–187 Deep Internal Processes and Continental Rifting. [10.1016/0040-1951\(87\)90262-9](http://dx.doi.org/10.1016/0040-1951(87)90262-9).
- Riis, F., Vollset, J., Sand, M., 1986. Tectonic Development of the Western Margin of the Barents Sea and Adjacent Areas. 131 pp. 661–675.
- Ryan, W.B.F., Carbotte, S.M., Coplan, J.O., O'Hara, S., Melkonian, A., Arko, R., Weissel, R.A., Ferrini, V., Goodwillie, A., Nitsche, F., Bonczkowski, J., Zemsky, R., 2009. Global multi-resolution topography synthesis: global multi-resolution topography synthesis. *Geochemistry, Geophysics, Geosystems* 10. <http://dx.doi.org/10.1029/2008GC002332> (n/a–n/a).
- Sapin, F., Ringenbach, J.-C., Rives, T., Pubellier, M., 2012. Counter-regional normal faults in shale-dominated deltas: origin, mechanism and evolution. *Marine and Petroleum Geology* 37:121–128. <http://dx.doi.org/10.1016/j.marpetgeo.2012.05.001>.
- Savva, D., Meresse, F., Pubellier, M., Chamot-Rooke, N., Lavier, L., Po, K.W., Franke, D., Steuer, S., Sapin, F., Auxietre, J.L., Lamy, G., 2013. Seismic evidence of hyper-stretched crust and mantle exhumation offshore Vietnam. *Tectonophysics* 608:72–83. <http://dx.doi.org/10.1016/j.tecto.2013.07.010>.
- Savva, D., Pubellier, M., Franke, D., Chamot-Rooke, N., Meresse, F., Steuer, S., Auxietre, J.L., 2014. Different expressions of rifting on the South China Sea margins. *Marine and Petroleum Geology* 58 (Part B):579–598 Evolution, Structure, and Sedimentary Record of the South China Sea and Adjacent Basins. [10.1016/j.marpetgeo.2014.05.023](http://dx.doi.org/10.1016/j.marpetgeo.2014.05.023).
- Schlüter, H.U., Hinz, K., Block, M., 1996. Tectono-stratigraphic terranes and detachment faulting of the South China Sea and Sulu Sea. *Marine Geology* 130:39–78. [http://dx.doi.org/10.1016/0025-3227\(95\)00137-9](http://dx.doi.org/10.1016/0025-3227(95)00137-9).
- Sibuet, J.-C., 1992. New constraints on the formation of the non-volcanic continental Galicia–Flemish Cap conjugate margins. *Journal of the Geological Society* 149:829–840. <http://dx.doi.org/10.1144/gsjgs.149.5.0829>.
- Skogseid, J., 2014. Extension on rifted continental margins: observation vs. models. *Geophysical Research Abstracts*. vol. 16. EGU General Assembly 2014. <http://meetingorganizer.copernicus.org/EGU2014/EGU2014-14913.pdf>.
- Skogseid, J., Pedersen, T., Eldholm, O., Larsen, B.T., 1992. Tectonism and magmatism during NE Atlantic continental break-up: the Vøring Margin. Geological Society of London, Special Publication 68:305–320. <http://dx.doi.org/10.1144/GSLSP.1992.068.01.19>.
- Soto, M., Morales, E., Veroslavsky, G., de Santa Ana, H., Ucha, N., Rodríguez, P., 2011. The continental margin of Uruguay: crustal architecture and segmentation. *Marine and Petroleum Geology* 28:1676–1689. <http://dx.doi.org/10.1016/j.marpetgeo.2011.07.001>.
- Stica, J.M., Zalán, P.V., Ferrari, A.L., 2014. The evolution of rifting on the volcanic margin of the Pelotas Basin and the contextualization of the Paraná–Etendeka LLP in the separation of Gondwana in the South Atlantic. *Marine and Petroleum Geology* 50:1–21. <http://dx.doi.org/10.1016/j.marpetgeo.2013.10.015>.
- Talwani, M., Eldholm, O., 1977. Evolution of the Norwegian-Greenland Sea. Geological Society of America Bulletin 88:969–999. [http://dx.doi.org/10.1130/0016-7606\(1977\)88<969:EOTNS>2.0.CO;2](http://dx.doi.org/10.1130/0016-7606(1977)88<969:EOTNS>2.0.CO;2).
- Taylor, B., Hayes, D.E., 1980. The tectonic evolution of the South China Basin. In: Hayes, D.E. (Ed.), *The Tectonic and Geologic Evolution of Southeast Asian Seas and Islands*. American Geophysical Union, pp. 89–104.
- Taylor, B., Hayes, D.E., 1983. Origin and history of the South China Sea Basin. In: Hayes, D.E. (Ed.), *The Tectonic and Geologic Evolution of Southeast Asian Seas and Islands: Part 2*. American Geophysical Union, pp. 23–56.
- Teisserenc, P., Villemin, J., 1990. Sedimentary basin of Gabon; geology and oil systems. *AAPG Memoir* 48, 117–199.
- Teixell, A., Labaume, P., Lagabrielle, Y., 2016. The crustal evolution of the west-central Pyrenees revisited: inferences from a new kinematic scenario. *Comptes Rendus Geosciences* 348:257–267 From Rifting to Mountain Building: The Pyrenean Belt. [10.1016/j.crte.2015.10.010](http://dx.doi.org/10.1016/j.crte.2015.10.010).
- Theissen, S., Rüpke, L.H., 2010. Feedbacks of sedimentation on crustal heat flow: new insights from the Vøring Basin, Norwegian Sea. *Basin Research* 22:976–990. <http://dx.doi.org/10.1111/j.1365-2117.2009.00437.x>.
- Tsikalas, F., Faleide, J.J., Kuszniir, N.J., 2008. Along-strike variations in rifted margin crustal architecture and lithosphere thinning between northern Vøring and Lofoten margin segments off mid-Norway. *Tectonophysics* 458:68–81 Geodynamics of Lithospheric Extension. [10.1016/j.tecto.2008.03.001](http://dx.doi.org/10.1016/j.tecto.2008.03.001).
- Unternehr, P., Péron-Pinvidic, G., Manatschal, G., Sutra, E., 2010. Hyper-extended crust in the South Atlantic: in search of a model. *Petroleum Geoscience* 16:207–215. <http://dx.doi.org/10.1144/1354-079309-904>.
- Van Avendonk, H.J.A., Lavier, L.L., Shillington, D.J., Manatschal, G., 2009. Extension of continental crust at the margin of the eastern Grand Banks, Newfoundland. *Tectonophysics* 468:131–148 Role of Magmatism in Continental Lithosphere Extension. [10.1016/j.tecto.2008.05.030](http://dx.doi.org/10.1016/j.tecto.2008.05.030).
- Vendeville, B., 1987. Champs de failles et tectonique en extension: Modélisation expérimentale. (phd thesis). Université Rennes 1.
- Wang, Y., Fan, W., Zhang, G., Zhang, Y., 2013. Phanerozoic tectonics of the South China Block: key observations and controversies. *Gondwana Research* 23:1273–1305. <http://dx.doi.org/10.1016/j.gr.2012.02.019>.
- Watts, A.B., Stewart, J., 1998. Gravity anomalies and segmentation of the continental margin offshore West Africa. *Earth and Planetary Science Letters* 156:239–252. [http://dx.doi.org/10.1016/S0012-821X\(98\)00018-1](http://dx.doi.org/10.1016/S0012-821X(98)00018-1).
- White, R., McKenzie, D., 1989. Magmatism at rift zones: the generation of volcanic continental margins and flood basalts. *Journal of Geophysical Research - Solid Earth* 94: 7685–7729. <http://dx.doi.org/10.1029/JB094iB06p07685>.

- White, R.S., Spence, G.D., Fowler, S.R., McKenzie, D.P., Westbrook, G.K., Bowen, A.N., 1987. Magmatism at rifted continental margins. *Nature* 330:439–444. <http://dx.doi.org/10.1038/330439a0>.
- Whitmarsh, R.B., Sawyer, D.S., Klaus, A., Masson, D.G. (Eds.), 1996. *Proceedings of the Ocean Drilling Program, 149 Scientific Results. Ocean Drilling Program*.
- Xiao, H.-B., Dahlen, F.A., Suppe, J., 1991. Mechanics of extensional wedges. *Journal of Geophysical Research - Solid Earth* 1978–2012:96. <http://dx.doi.org/10.1029/91JB00222>.
- Yin, A., 1994. Mechanics of wedge-shaped fault blocks: 2. An elastic solution for extensional wedges. *Journal of Geophysical Research - Solid Earth* 99:7045–7055. <http://dx.doi.org/10.1029/93JB02389>.
- Ziegler, P.A., 1983. Crustal thinning and subsidence in the North Sea. *Nature* 304:561. <http://dx.doi.org/10.1038/304561a0>.



**Laurent Jolivet** is professor of the University of Orléans (France). He received his Ph.D. from Université Pierre et Marie Curie in Paris. He has developed research in tectonics and geodynamics, focussing on upper plate deformation in subduction zones, back-arc basins opening, exhumation of HP-LT metamorphic rocks, lithosphere-asthenosphere coupling, metamorphic core-complexes and detachments, dynamics of extension, Mediterranean tectonics and geodynamics, tectonic evolution of East Asia. He is the co-PI of the ERC advanced research grant RHEOLITH together with E. Burov, a project aimed at a better understanding of strain localisation in the continental lithosphere.



**Camille Clerc** is associate professor at the University of New Caledonia. He earned his Ph.D. in structural geology in 2012 at the Université Pierre et Marie Curie in Paris for studying the aborted cretaceous rift system fossilized in the Pyrenees. Since 2009, Camille is studying the architecture and thermal evolution of passive margins, both from fossil margins exhumed in orogens (Pyrenees) and from present day passive margins. Recruited at the University of New Caledonia in 2016, he now explores the extension of the drawn continent Zealandia and investigates the geological setting of the New Caledonian ophiolite.



**Jean-François Ballard** earned his Ph.D. in structural geology in 1989 from Rennes University, France under direction of Prof. Jean-Pierre Brun (analog modeling of compressive and extensional systems, field works in Brittany, France). In 1989, JF started working at IFP and as an external consultant for Elf, which he joined in 1991. Geologist in the structural group he did field works in Argentina, Venezuela, Peru and structural studies on various tectonic context from margin to fold-and-thrust belts. From 2001 to 2004, he integrated Total E&P Bolivie as an exploration geologist. Since 2005, JF is based in Pau and mainly dedicates himself to structural support to Total affiliates (Bolivia, Nigeria, Gabon,...) and headquarters group (new ventures etc...).



**Jean-Claude Ringenbach** studied in Ecole Normale Supérieure and he earned his Ph.D. in structural geology and geodynamics in 1992 from Nice University, France. He worked for Elf and then Total since 1992. Exploration Geologist in Total Affiliates, he occupied several positions: basin geologist and deep offshore Exploration Manager (1997–2001) and Chief Geologist in Libya (2001–2004). From 2004 to 2007 JC took over the Fold-and-Thrust Belts thematic in the New Venture Group in Paris. From 2007 to 2013 he was the Head of the Structural Geology Group in Pau and has been appointed Total's Expert in Structural Geology in 2013. JC's experience encompasses various fields of basin structural geology and is based on a large worldwide geological knowledge. His expertise is supported by academic projects on Fold-and-Thrust Belts, Salt Tectonics and Continental Margins.